

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number  
**WO 01/36464 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 14/00**

(21) International Application Number: **PCT/US00/31187**

(22) International Filing Date:  
13 November 2000 (13.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/166,128 18 November 1999 (18.11.1999) US  
60/201,837 3 May 2000 (03.05.2000) US

(71) Applicants (for all designated States except US): **PIONEER HI-BRED INTERNATIONAL, INC.** [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50309 (US). **CURAGEN CORPORATION** [US/US]; 555 Long Wharf Drive, 13th Floor, New Haven, CT 06511 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BIDNEY, Dennis, L.** [US/US]; 8385 Plum Drive, Urbandale, IA 50322 (US).

**DUVICK, Jonathan, P.** [US/US]; 1707 - 38th Street, Des Moines, IA 50310 (US). **HENDRICK, Carol** [US/US]; 1707 - 38th Street, Des Moines, IA 50310 (US). **HU, Xu** [CA/US]; 4700 - 103rd Street, Urbandale, IA 50322 (US). **LU, Guihua** [CN/US]; 4600 - 62nd Street, Urbandale, IA 50322 (US). **CRASTA, Oswald, R.** [IN/US]; 95-2C Florence Road, Branford, CT 06405 (US).


(74) Agents: **WILLIAMSON, Kelly, J. et al.**; Alston & Bird LLP, P.O. Drawer 34009, Charlotte, NC 28234-4009 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European

[Continued on next page]

(54) Title: **SUNFLOWER rhoGAP, LOX, ADH, AND SCIP-1 POLYNUCLEOTIDES AND METHODS OF USE**



(57) Abstract: Methods and compositions for modulating development and defense response are provided. Nucleotide sequences encoding a sunflower rhoGAP, LOX, ADH, and SCIP-1 protein are provided. Nucleotide sequences comprising the LOX and SCIP-1 promoters are also provided. The sequences can be used in expression cassettes for modulating development, developmental pathways, and the plant defense response. Transformed plants, plant cells, tissues, and seed are also provided.



**WO 01/36464 A2**



patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

— *Without international search report and to be republished upon receipt of that report.*

## SUNFLOWER RhoGAP, LOX, ADH, AND SCIP-1 POLYNUCLEOTIDES AND METHODS OF USE

### FIELD OF THE INVENTION

The invention relates to the field of the genetic manipulation of plants, particularly the modulation of gene activity and development in plants and increased disease resistance.

### 5 BACKGROUND OF THE INVENTION

Disease in plants is caused by biotic and abiotic causes. Biotic causes include fungi, viruses, bacteria, and nematodes. An example of the importance of plant disease is illustrated by phytopathogenic fungi, which cause significant annual crop yield losses as well as devastating epidemics. Plant disease outbreaks have resulted in catastrophic crop failures that have triggered famines and caused major social change. Pathogenic fungi attack all of the approximately 300,000 species of flowering plants, however, a single plant species can be host to only a few fungal species, and similarly, most fungi usually have a limited host range. Generally, the best strategy for plant disease control is to use resistant cultivars selected or developed by plant breeders for this purpose. However, the potential for serious crop disease epidemics persists today, as evidenced by outbreaks of the Victoria blight of oats and southern corn leaf blight. Molecular methods of crop protection have the potential to implement novel mechanisms for disease resistance and can also be implemented more quickly than traditional breeding methods. Accordingly, molecular methods are needed to supplement traditional breeding methods to protect plants from pathogen attack.

A host of cellular processes enable plants to defend themselves against disease caused by pathogenic agents. These defense mechanisms are activated by initial pathogen infection in a process known as elicitation. In elicitation, the host plant recognizes a pathogen-derived compound known as an elicitor; the plant then activates disease gene expression to limit further spread of the invading microorganism. It is generally believed that to overcome these plant defense mechanisms, plant pathogens must find a way to suppress elicitation as well as to

overcome more physically-based barriers to infection, such as reinforcement and/or rearrangement of the actin filament networks near the cell's plasma membrane.

Thus, the present invention solves needs for enhancement of the plant's defensive elicitation response via a molecularly based mechanism that can be quickly  
5 incorporated into commercial crops.

### SUMMARY OF THE INVENTION

The present invention provides nucleotide sequences that may find use in modulating development, developmental pathways, and the plant pathogen defense  
10 system. Particularly, the nucleotide and amino acid sequences for a sunflower rhoGTPase-Activating Protein (rhoGAP), Lipoxxygenase (LOX), Alcohol Dehydrogenase (ADH), and *Sclerotinia*-Inducible Protein-1 (SCIP-1) are provided.

In particular, the methods and compositions can be used to modulate plant development. More specifically, methods and compositions of the invention may be  
15 used for enhancing resistance to plant pathogens including fungal pathogens, plant viruses, and the like. The method involves stably transforming a plant with a nucleotide sequence capable of modulating the plant pathogen defense system operably linked with a promoter capable of driving expression of a gene in a plant cell. The disease resistance genes of the present invention additionally find use in  
20 manipulating these processes in transformed plants and plant cells.

Transformed plants, plant cells, and seeds, as well as methods for making such plants, plant cells, and seeds are additionally provided. It is recognized that a variety of promoters will be useful in the invention, the choice of which will depend in part upon the desired level of expression of the disclosed nucleotide sequences. It is  
25 recognized that the levels of expression can be controlled to modulate the levels of expression in the plant cell.

Methods and compositions for regulating gene expression in a plant are also provided. Novel nucleotide sequences for inducible plant promoters derived from the LOX and SCIP-1 genes are provided. The methods comprise transforming a plant  
30 with a nucleotide sequence of interest operably linked to the LOX or SCIP-1 promoters. Exposure of the transformed plant to a stimulus activates, within the exposed tissue of the plant, transcription of the nucleotide sequence of interest.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 schematically illustrates an expression vector containing the ubiquitin promoter operably linked the rhoGAP, LOX, ADH, and SCIP-1 nucleotide sequences.

Figure 2 schematically illustrates an expression vector used for plant transformation containing the LOX or SCIP-1 promoter nucleotide sequences operably linked to the nucleotide sequences encoding the GUS reporter protein.

## DETAILED DESCRIPTION OF THE INVENTION

Overview

The present invention provides, inter alia, compositions and methods for modulating the total level of proteins of the present invention and/or altering their ratios in a plant. By "modulation" is intended an increase or a decrease in a particular character, quality, substance, or response.

The compositions comprise sunflower nucleic acid and amino acid sequences. Particularly, the nucleotide and amino acid sequence for a sunflower rhoGAP (SEQ ID NOS: 1 and 2), LOX (SEQ ID NOS: 3 and 4), ADH (SEQ ID NOS: 6 and 7) and SCIP-1 (SEQ ID NOS: 8 and 9) are provided. As discussed in more detail below, the sequences of the invention are involved in many basic biochemical pathways that regulate plant growth, development, and pathogen resistance. Methods are provided for the expression of these sequences in a host plant to modulate plant development, developmental pathways, and defense responses. The method involves stably transforming a plant with a nucleotide sequence capable of modulating the plant pathogen defense system operably linked with a promoter capable of driving expression of the nucleotide sequence in a plant cell.

Also provided are LOX and SCIP-1 promoter sequences set forth in SEQ ID NO: 5 and SEQ ID NO: 10, respectively. Methods are provided for the regulated expression of a nucleotide sequence of interest that is operably linked to the LOX or SCIP-1 promoter sequences disclosed herein. Nucleotide sequences operably linked to the LOX or SCIP-1 promoters are transformed into a plant cell. Exposure of the transformed plant to a stimulus, induces transcriptional activation of the nucleotide sequences operably linked to the LOX or SCIP-1 promoters.

*rhoGTPase-activating protein (rhoGAP)*

A nucleic acid sequence encoding a rhoGAP polypeptide from sunflower is provided. The rhoGAP sequence shares homology to the conserved rhoGAP genes from humans. The sunflower rhoGAP amino acid sequence shares about 40%  
5 homology with the human p50rhoGAP sequence and about 80% homology with an *Arabidopsis* hypothetical 23.6 kDa protein.

RhoGAPs are a central part of an evolutionarily conserved regulatory system that are involved in cell growth and differentiation. Thus the sequence of the invention finds use in controlling or modulating cell division, differentiation,  
10 development, as well as the defense response. Transformed plants can be obtained having altered metabolic states with respect to cell division and cellular processes as well as development and defense response; hence, the methods and compositions may find uses in affecting or studying differentiation.

RhoGAP proteins have been shown to interact with rho members of the ras  
15 superfamily. Ras oncogenes were initially found to play an important role in human cancers and have since been shown to play important roles in regulation of cell growth and differentiation. Further, the rhoGAP proteins affect the activity of rhoGTPases (also called rho proteins), which act as molecular switches to regulate affected processes. The rho family of "G proteins" have a GTP-bound form and a  
20 GDP-bound form; the relative amount of the GDP-bound form is increased by GTPase activating proteins, or GAPs, which stimulate the intrinsic GTPase activity of the rho proteins.

Processes affected by GAPs include the transduction of hormone signals across cell plasma membranes and the regulation of intracellular transport pathways.  
25 For example, rhoGTP-binding proteins have been shown to control signal transduction pathways connecting the activation of actin polymerization to activation of cellular growth factor receptors. Hence, the compositions and methods of the invention find use in the activation or modulation of the cellular actin cytoskeleton. Although there is a great deal of conservation among members of the rhoGAP family,  
30 there are a large number of different proteins that contain the rhoGAP domain, and many of these proteins are large and multifunctional. Thus, the rhoGAP genes and/or proteins may contain different elements or motifs or sequence patterns that modulate or affect the activity, subcellular localization, and/or target of the rhoGAP protein.

Such elements, motifs, or sequence patterns may be useful in engineering novel enzymes for reducing or enhancing gene expression in particular tissues.

RhoGAP proteins activate rho genes and the related rac genes, which both stimulate actin polymerization. The rho proteins in mammalian systems have been shown to regulate the formation of multi-molecular complexes that are associated with polymerized actin located at the plasma membrane of the cell. Such complexes include actin stress fibers and focal adhesions in fibroblasts as well as the actin-driven phenomenon called membrane ruffling, which is exhibited by many cell types in response to extracellular stimuli. Rho proteins have also been shown to play roles in epithelial cell migration in response to wounding. Expression of the sequences of the invention can be used to modulate or regulate the expression of corresponding GTP-binding proteins, i.e., rho, rac, etc. Hence, the compositions and methods of the invention find use in the activation or modulation of the cellular actin cytoskeleton and other actin-based structures and actin-related processes.

The RhoGAP gene of the present invention additionally finds use in enhancing the plant pathogen defense system. Early plant-cell defense responses include the rearrangement of the cellular actin cytoskeleton to protect the cell from attack. RhoGAP genes are involved in cellular signaling cascades such as the oxidative burst that comprises part of the early defense response in plants. Hence, the compositions and methods of the invention can be used for enhancing resistance to plant pathogens including fungal pathogens, plant viruses, and the like.

#### *Lipoxygenase*

A nucleic acid sequence encoding a LOX polypeptide from sunflower is also provided. The sunflower LOX polypeptide shares homology with other known LOX proteins from potato, tomato, cowpea, *Arabidopsis*, and rice.

The LOX protein has been implicated in a number of important plant developmental processes. LOX catalyzes the hydroperoxidation of polyunsaturated fatty acids containing *cis*, *cis*-1,4-pentadiene-conjugated double-bonds. The primary products of LOX-catalyzed reactions are fatty acid hydroperoxides, which are typically metabolized into molecules with known or suspected regulatory activity. For example, LOX derived fatty acid hydroperoxides are precursors to traumatin and jasmonic acid. Traumatin induces cell division and may be involved in the plant

wounding response (Zimmerman *et al.* (1979) *Plant Physiol.* 63:536-541).

Jasmonates have been implicated as signal transduction molecules in the response of plants to stress, particularly wounding and pathogen attack (Farmer *et al.* (1992) *Cell* 4:129-134). Therefore, the sunflower LOX gene may play an important role in cell  
5 division and defense signal transduction pathways that are regulated by the biosynthesis of traumatin and jasmonic acid.

The LOX gene has also been implicated in the regulation of coordinated gene activation in response to wounding. It is speculated that resistance to pathogen attack is the result of the coordinated accumulation of secondary metabolites and protein  
10 products. Some of these products, such as proteinase inhibitors, may directly interfere with digestibility of the injected tissue whereas others products may affect food intake. A potato LOX gene, 13-LOX, has been shown to control the expression levels of proteinase inhibitors in a wounding response to insect feeding (Royo *et al.* (1999) *Proc. Natl. Acad. Sci. USA* 96:1146-1151). Therefore, the LOX sequences of the  
15 present invention may find use in an antifeedant strategy by regulating proteinase inhibitor levels in plants, and thereby controlling insect and nematode pathogens.

Additionally, LOX-derived fatty acid hydroperoxides and free radical species are cytotoxic and are capable of damaging membranes, proteins, and DNA (Hildebrand *et al.* (1998) *Curr. Top. Plant Biochem. Physiol.* 7:201-219). Therefore,  
20 LOXs may play a role in membrane degradation observed during senescence, wounding, and the hypersensitive response to pathogen attack.

LOX proteins may also play an important role in plant growth and development. There is a positive correlation between LOX activity levels within an organ and its rate of elongation. The concomitant increase in LOXs and the enzymes  
25 involved in the metabolism of LOX-derived fatty hydroperoxides is consistent with a role for LOX in generating lipid-derived growth regulators.

Furthermore, in plants, the LOX proteins may be involved in lipid turnover and fat mobilization. Hence, the compositions and methods of the invention find use the turnover of lipids in the developing seedling.

30 Hence, the LOX sequences of the present invention may be used to modulate many important developmental processes, such as, cell division, seed germination, plant growth and senescence, and/or to enhanced plant resistance to environmental stresses, such as, wounding and pathogen attacks.



The present invention also provides the nucleotide sequences of the LOX promoter. The promoter sequence contains *cis*-elements that resemble W-boxes, TTGACC (nucleotides 322-327 of SEQ ID NO: 5), and G-boxes, CACGTG (nucleotides 722-727 of SEQ ID NO: 5).

5           G-box and G-box like sequences are involved in the regulation of a variety of unrelated genes and activate transcription in response to various stimuli including: exposure to visible and UV light (Chattopadhyay *et al.* (1998) *Plant Cell* 10:673-683), dehydration-stress (Lam *et al.* (1991) *J Biol Chem* 266: 17131-17135 and Dolferus *et al.* (1994) *Plant Physiol* 105: 1075-1087), cold stress (Dolferus *et al.* (1994) *Plant*  
10   *Physiol* 105: 1075-1087), abscisic acid (Marcotte *et al.* (1989) *Plant Cell* 1: 969-976), sucrose (Urwin *et al.* (1997) *Plant Mol Biol* 35:929-942), and plant pathogen defense response (Wolfgang *et al.* (1997) *EMBO Journal* 16:726-738). In addition, G-box-like sequences were also found to determine tissue-preferred expression patterns (Salinas *et al.* (1992) *Plant Cell* 4: 1485-1493 and Thomas (1993) *Plant Cell* 5: 1401-  
15   1410). Functional analysis of G-box containing promoters has shown that the nucleotide sequences immediately flanking the G-box and/or additional *cis*-acting promoter elements are often required for the G-box to influence transcription activation.

          The W-box promoter elements are involved in elicitor-induced gene  
20   expression. W-boxes have been identified in a several plant promoters including, for example, members of the WRKY family (Eulgem *et al.* (1999) *EMBO J.* 18:4689-4699) and members of the pathogenesis-related protein family (Rushton *et al.* (1996) *EMBO J.* 15:5690-5700). The fungal elicitor responsiveness of these genes is mediated mainly by the presence of the W-boxes in the promoter elements.

25           Hence, the LOX promoter sequences find use in the regulated expression of an operably linked heterologous gene of interest. More specifically, the nucleotide sequence may find use as an inducible promoter, more specifically, a pathogen-inducible promoter.

### 30   *Alcohol Dehydrogenase*

A nucleic acid sequence encoding an ADH protein from sunflower is also provided. Sequences of the sunflower alcohol dehydrogenase protein (ADH) share about 85-95% sequence homology with plant alcohol dehydrogenases from garden

lettuce, potato, tomato, apple, and maize. ADH is an important enzyme in anaerobic metabolism, and it is usually encoded by a small multigene family in flowering plants. In both maize and *Arabidopsis*, the gene is expressed in seeds, roots, and pollen grains, whereas green aerial plant parts are devoid of detectable ADH activity.

- 5 ADH has been implicated in responses to a number of environmental stresses, including low oxygen, drought, salinity, cold acclimation, freezing tolerance, flooding, and wounding. See, for example, Zeevaart *et al.* (1988) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 39:439-473; Sanchez *et al.* (1991) *Abscisic Acid, Physiology and Biochemistry*, Bio Scientific Publishers, Oxford, UK, pp. 210-216; and Bruxelles  
10 *et al.* (1996) *Plant Physiol.* 111:381-391. The ADH sequence of the present invention may find use in modulating a plant's response to adverse environmental stresses.

- The sunflower ADH sequences may also find use in modulating the developmental process of fruit ripening. ADH reduces aldehydes to alcohols. Modulation of ADH levels in ripening fruit has been shown to influence the balance  
15 between some of the aldehydes and the corresponding alcohols associated with flavor production. Hence, the compositions and methods of the present invention may find use in the modulation of ADH protein levels leading to a more intense "ripe fruit" flavor. See, for example, Speirs *et al.* (1998) *Plant Physiol.* 117:1047-1058.

- The ADH sequences of the invention may additionally find use in enhancing  
20 plants defense response. Under low oxygen conditions, i.e., a hypoxic state, ADH plays a crucial role in cell survival. ADH serves as the major terminal dehydrogenase in regenerating oxidizing power in mature roots, thereby allowing glycolysis to continue in the absence of oxygen. Treatment of a plant with an elicitor increases the levels of active oxygen species in the plant cells and leads to a transient state of  
25 oxidative stress. See, for example, Robertson *et al.* (1995) *Plant Molecular Biology* 27:59-67. Since aerobic respiration is compromised as a result of elicitor action, the ADH sequences of the present invention may find use in modulating a plant's defense against pathogens.

30 *Sclerotinia-inducible protein-1 (SCIP-1)*

The nucleic acid sequence encoding a novel sunflower protein, designated *Sclerotinia*-Inducible Protein-1, SCIP-1, is also provided. SCIP-1 has limited homology with hypothetical proteins from several bacteria.

Transcript levels of SCIP-1 increase in both lesion mimic transgenic plants and *Sclerotinia*-infected plants. The accumulation of SCIP-1 in lesion mimic and infected sunflower plants implicates that the protein is involved in the plant defense response to *Sclerotinia* and other pathogens. Hence, the compositions and methods of the invention may find uses for enhancing resistance to plant pathogens, including fungal pathogens, plant viruses, and the like.

Furthermore, a PSI-Blast search revealed that the SCIP-1 sequence of the invention fall into a class of flowering-related plant proteins (CEN and others), as well as some phosphatidylethanolamine-binding proteins (PEBP). The CEN-related proteins are known to be related to a class of phosphatidylethanolamine-binding proteins (PEBP). Banfield *et al.* ((2000) *J Mol Biol* 297:1159-70) determined the crystal structure of the centroradialis protein from *Antirrhinum*. The structure confirmed what had been suspected by sequence homology studies: that the CEN plant proteins are a subset of the family of PEBPs. Mammalian forms of PEBP are involved in inhibition of MAP kinase signaling, which is a central signaling cascade regulating cell differentiation (Banfield *et al.* (2000) *J Mol Biol* 297:1159-70). The structure of these proteins suggests that they may play a role in membrane signal transduction (Banfield *et al.* (1998) *Structure* 6:1245-54). In addition, another recent study (Kuramitsu *et al.* (2000) *Electrophoresis* 21:660-4) showed that a line of mammalian cells resistant to tumor necrosis factor-alpha contained elevated levels of a protein identified as a PEBP. The report suggested that this PEBP could be responsible for the resistance of certain cell lines to tumor necrosis factor induced cell death. Hence, the SCIP-1 polypeptide of the invention may play a role in signaling, membrane transduction, or in the regulation of cell death.

Furthermore, flowering plants exhibit two types of inflorescence architecture: determinate and indeterminate. The centroradialis mutation causes the normally indeterminate inflorescence of *Antirrhinum* to terminate in a flower. CEN-related protein have therefore been shown to influence maintenance of the indeterminate state of inflorescence meristems (Pnueli *et al.* (1998) *Development* 125:1979-1989; Bradley *et al.* (1996) *Nature* 379:791-7; Bradley-Desmond *et al.* (1997) *Science* 275:80-83; and Amaya *et al.* (1999) *Plant Cell* 11:1405-1417). In addition, the SCIP-1 shares homology to Terminal Flower 1 (TFL1) from both *Arabidopsis thaliana* and *Brassica*. TFL1 has also been shown to influence inflorescence meristem identity.

See, for example, Mimida-Naozumi *et al.* (1999) *Plant-Science* 142: 155-162 and Ohshima *et al.* (1997) *Mol Gen Genet* 254:186-94. Hence, the SCIP-1 sequences of the invention find use in influencing the state of inflorescence of meristem development.

- 5           The present invention also provides the nucleotide sequences of the SCIP-1 promoter. The promoter sequence contains *cis*-elements that resemble W-boxes, GTCAA (nucleotides 364-368 and 371-375 of SEQ ID NOS: 8 and 10), and G-boxes, CACGTG (nucleotides 415-420 of SEQ ID NOS: 8 and 10). As with the LOX promoter sequences, the SCIP-1 promoter sequences may find use in the regulated  
10 expression of an operably linked heterologous gene of interest. More specifically, the nucleotide sequence may find use as an inducible promoter, more specifically, a pathogen-inducible promoter.

#### Compositions

- 15           Compositions of the invention include the polynucleotide sequences of the sunflower rhoGAP, LOX, ADH, and SCIP-1 genes. In addition, the LOX and SCIP-1 promoter nucleotide sequences are also provided. The polypeptides encoded by those sequences may be involved in various plant developmental processes, including the plant pathogen defense response.
- 20           In particular, the present invention provides for isolated nucleic acid molecules comprising nucleotide sequences encoding the amino acid sequences shown in SEQ ID NOS: 2, 4, 7, and 9 or the nucleotide sequences encoding the DNA sequences deposited in a bacterial host as Patent Deposit Nos. PTA-284 and PTA-285 (DNA sequences corresponding to rhoGAP), PTA-286 (DNA sequences  
25 corresponding to ADH), PTA-287 (DNA sequences corresponding to LOX), PTA-288 (DNA sequences corresponding to SCIP-1), or the DNA sequences obtained from the overlapping clones deposited in a bacterial host as Patent Deposit Nos. PTA-284 and PTA-285. Further provided are polypeptides having an amino acid sequence encoded by a nucleic acid molecule described herein, for example those set forth in  
30 SEQ ID NOS: 1, 3, 6, and 8 or those deposited in a bacterial host as Patent Deposit Nos. PTA-284, PTA-285, PTA-286, PTA-287, PTA-288 and fragments and variants thereof.

The present invention further provides for isolated nucleic acid molecules comprising nucleotide sequences shown in SEQ ID NO: 5 and SEQ ID NO: 10, or nucleotide sequences encoding the DNA sequences deposited in a bacterial host as Patent Deposit Nos. PTA-559 and PTA-1721, and fragments and variants thereof.

5 By "DNA sequences obtained from the overlapping clones" is intended that the complete DNA sequence of the rhoGAP sequence of the invention (SEQ ID NO: 1) can be obtained by sequencing the two individual clones that together comprise the entire rhoGAP sequence.

Plasmids containing the nucleotide sequences of the invention were deposited  
10 with the Patent Depository of the American Type Culture Collection (ATCC), Manassas, Virginia, and assigned Accession Nos. PTA-284, PTA-285, PTA-286, PTA-287, PTA-288, PTA-559, and PTA-1721. Plasmids having the Accession Nos. PTA-284, PTA-285, PTA-286, PTA-287, and PTA-288 were deposited on June 30, 1999. The plasmid having Accession No. PTA-559 was deposited on August 20,  
15 1999, and the plasmid deposited as Accession No. PTA-1721 was deposited on April 26, 2000. Two of these plasmids, designated as Accession No. PTA-284 and Accession No. PTA-285, contained overlapping clones. The plasmids deposited as PTA-284 and PTA-285 comprise the 5' and the 3' end of the rhoGAP sequence, respectively. It is noted, however, that clones deposited as PTA-284 and PTA-285  
20 contain common sequences at the regions where they overlap. One of skill in the art by sequencing the clones and aligning the overlap may obtain the entire sequence of the sunflower rhoGAP. These deposits will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. These deposits were made merely as a  
25 convenience for those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112.

The invention encompasses isolated or substantially purified nucleic acid or protein compositions. An "isolated" or "purified" nucleic acid molecule or protein, or biologically active portion thereof, is substantially free of other cellular material, or  
30 culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Preferably, an "isolated" nucleic acid is free of sequences (preferably protein encoding sequences) that naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the

nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell  
5 from which the nucleic acid is derived. A protein that is substantially free of cellular material includes preparations of protein having less than about 30%, 20%, 10%, 5%, (by dry weight) of contaminating protein. When the protein of the invention or biologically active portion thereof is recombinantly produced, preferably culture medium represents less than about 30%, 20%, 10%, or 5% (by dry weight) of  
10 chemical precursors or non-protein-of-interest chemicals.

Fragments and variants of the disclosed nucleotide sequences and proteins encoded thereby are also encompassed by the present invention. By "fragment" is intended a portion of the nucleotide sequence or a portion of the amino acid sequence and hence protein encoded thereby. Fragments of a nucleotide sequence may encode  
15 protein fragments that retain the biological activity of the native protein and hence affect development, developmental pathways, and defense response by retaining rhoGAP-, LOX-, ADH-, or SCIP-1-like activity. Alternatively, fragments of a nucleotide sequence that are useful as hybridization probes generally do not encode  
20 fragment proteins retaining biological activity. Thus, fragments of a nucleotide sequence may range from at least about 20 nucleotides, about 50 nucleotides, about 100 nucleotides, and up to the full-length nucleotide sequence encoding the proteins of the invention.

A fragment of a rhoGAP nucleotide sequence that encodes a biologically active portion of a rhoGAP protein of the invention will encode at least 12, 25, 30, 50,  
25 100, 150, or 200 contiguous amino acids, or up to the total number of amino acids present in a full-length rhoGAP protein of the invention (for example, 201 amino acids for SEQ ID NO: 2).

A fragment of a LOX nucleotide sequence that encodes a biologically active portion of a LOX protein of the invention will encode at least 22, 30, 50, 100, 150,  
30 200, 300, 400, 500, 600, 700, 800, 900 contiguous amino acids, or up to the total number of amino acids present in a full-length LOX protein of the invention (for example, 901 amino acids for SEQ ID NO: 3).

A fragment of an ADH nucleotide sequence that encodes a biologically active portion of an ADH protein of the invention will encode at least 92, 100, 150, 200, 250, 300, 350, 400 contiguous amino acids, or up to the total number of amino acids present in a full-length ADH protein of the invention (for example, 381 amino acids for SEQ ID NO: 7).

A fragment of a SCIP-1 nucleotide sequence that encodes a biologically active portion of a SCIP-1 protein of the invention will encode at least 8, 15, 25, 30, 50, 100, or 150 contiguous amino acids, or up to the total number of amino acids present in a full-length SCIP-1 protein of the invention (for example, 168 amino acids for SEQ ID NO: 9).

Fragments of a rhoGAP, LOX, ADH, and SCIP-1 nucleotide sequence that are useful as hybridization probes or PCR primers generally need not encode a biologically active portion of a rhoGAP, LOX, ADH, or SCIP-1 protein. Thus, a fragment of a rhoGAP, LOX, ADH, or SCIP-1 nucleotide sequence may encode a biologically active portion of a rhoGAP, LOX, ADH, or SCIP-1 protein, or it may be a fragment that can be used as a hybridization probe or PCR primer using methods disclosed below. A biologically active portion of a rhoGAP, LOX, ADH, or SCIP-1 protein can be prepared by isolating a portion of one of the rhoGAP, LOX, ADH, or SCIP-1 nucleotide sequences of the invention, expressing the encoded portion of the rhoGAP, LOX, ADH, or SCIP-1 protein (e.g., by recombinant expression *in vitro*), and assessing the activity of the encoded portion of the rhoGAP, LOX, ADH, or SCIP-1 protein. Nucleic acid molecules that are fragments of a rhoGAP, LOX, ADH, SCIP-1 nucleotide sequence comprise at least 16, 20, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, or 800 nucleotides, or up to the number of nucleotides present in a full-length rhoGAP, LOX, ADH, or SCIP-1 nucleotide sequence disclosed herein (for example, 824 nucleotides for SEQ ID NO: 1, 3806 nucleotides for SEQ ID NO: 3, 1403 nucleotides for SEQ ID NO: 6, and 746 nucleotide sequences for SEQ ID NO: 8).

By "variants" is intended substantially similar sequences. For nucleotide sequences, conservative variants include those sequences that, because of the degeneracy of the genetic code, encode the amino acid sequence of one of the polypeptides of the invention. Naturally occurring allelic variants such as these can be identified with the use of well-known molecular biology techniques, as, for

example, with polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant nucleotide sequences also include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis but which still encode a rhoGAP, LOX, ADH, or SCIP-1 protein of the invention. Generally, variants of a particular nucleotide sequence of the invention will have at least 40%, 50%, 60%, 70%, generally at least 75%, 80%, 85%, preferably about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, and more preferably about 98%, 99%, or more sequence identity to that particular nucleotide sequence as determined by sequence alignment programs described elsewhere herein using default parameters.

By "variant" protein is intended a protein derived from the native protein by deletion (so-called truncation) or addition of one or more amino acids to the N-terminal and/or C-terminal end of the native protein; deletion or addition of one or more amino acids at one or more sites in the native protein; or substitution of one or more amino acids at one or more sites in the native protein. Variant proteins encompassed by the present invention are biologically active, that is they continue to possess the desired biological activity of the native protein, hence they will continue to possess rhoGAP, LOX, ADH, or SCIP-1 activity. Such variants may result from, for example, genetic polymorphism or from human manipulation. Biologically active variants of a native rhoGAP, LOX, ADH, or SCIP-1 protein of the invention will have at least 40%, 50%, 60%, 70%, generally at least 75%, 80%, 85%, preferably about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to the amino acid sequence for the native protein as determined by sequence alignment programs described elsewhere herein using default parameters. A biologically active variant of a protein of the invention may differ from that protein by as few as 1-15 amino acid residues, as few as 1-10, such as 6-10, as few as 5, as few as 4, 3, 2, or even 1 amino acid residue.

Biological activity of the rhoGAP, LOX, ADH and SCIP-1 polypeptides can be assayed by any method known in the art. Assays to measure the developmental pathways and defense responses that are influenced by the rhoGAP, LOX, ADH and SCIP-1 polypeptides having rhoGAP-, LOX-, ADH-, and SCIP-1-like activity are well known in the art. Furthermore, assays to detect rhoGAP-like activity include GTP binding assays (Borg *et al.* (1994) *Plant Mol. Biol.* 27:175-187); interactions with Rac or Ras (Dieckman *et al.* (1995) *EMBO J.* 14:5297-5305 and Van Aelst *et al.*



(1996) *EMBO J.* 15:3778-3786); and GTPase and GTPase activating activity assays (Borg *et al.* (1999) *FEBS Letters* 453:341-345). Assays to detect LOX-like activity include, for example, assays to measure LOX enzymatic activity (Maach *et al.* (1997) *Plant Physiol.* 114:1561-1566, Royo *et al.* (1996) *J. Biol. Chem.* 35:21012-21019 and  
5 Voros *et al.* (1998) *FEBS Letters* 251:36-44). Assays to detect ADH-like activity include, for example, ADH enzymatic activity assays (Torres *et al.* (1976) *Biochem. Genetics* 14:87-97).

The proteins of the invention may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such  
10 manipulations are generally known in the art. For example, amino acid sequence variants of the rhoGAP, LOX, ADH, or SCIP-1 proteins can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods in Enzymol.* 154:367-382; US Patent No.  
15 4,873,192; Walker and Gastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff *et al.* (1978) *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington,  
20 D.C.), herein incorporated by reference. Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferred.

Thus, the genes and nucleotide sequences of the invention include both the naturally occurring sequences as well as mutant forms. Likewise, the proteins of the invention encompass both naturally occurring proteins as well as variations and  
25 modified forms thereof. Such variants will continue to possess the desired rhoGAP-, LOX-, ADH-, or SCIP-1-like activity. Obviously, the mutations that will be made in the DNA encoding the variant must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary  
30 mRNA structure. See, EP Patent Application Publication No. 75,444.

The deletions, insertions, and substitutions of the protein sequences encompassed herein are not expected to produce radical changes in the characteristics of the protein. However, when it is difficult to predict the exact effect of the

substitution, deletion, or insertion in advance of doing so, one skilled in the art will appreciate that the effect will be evaluated by routine screening assays. That is, the activity can be evaluated by either an enhanced resistance to pathogens or a modulation in a plant developmental process when expression of the protein sequence is altered.

Variant nucleotide sequences and proteins also encompass sequences and proteins derived from a mutagenic and recombinogenic procedure such as DNA shuffling. With such a procedure, one or more different rhoGAP, LOX, ADH, or SCIP-1 coding sequences can be manipulated to create a new rhoGAP, LOX, ADH, or SCIP-1 possessing the desired properties. In this manner, libraries of recombinant polynucleotides are generated from a population of related sequence polynucleotides comprising sequence regions that have substantial sequence identity and can be homologously recombined *in vitro* or *in vivo*. For example, using this approach, sequence motifs encoding a domain of interest may be shuffled between the rhoGAP, LOX, ADH, or SCIP-1 gene of the invention and other known rhoGAP, LOX, ADH, or SCIP-1 genes to obtain a new gene coding for a protein with an improved property of interest, such as an increased  $K_m$  in the case of an enzyme. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer (1994) *Nature* 370:389-391; Crameri *et al.* (1997) *Nature Biotech.* 15:436-438; Moore *et al.* (1997) *J. Mol. Biol.* 272:336-347; Zhang *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Crameri *et al.* (1998) *Nature* 391:288-291; and U.S. Patent Nos. 5,605,793 and 5,837,458.

The compositions of the invention also include isolated nucleic acid molecules comprising the promoter nucleotide sequences set forth in SEQ ID NO: 5 and SEQ ID NO: 10. By "promoter" is intended a regulatory region of DNA usually comprising a TATA box (nucleotides 808-901 of SEQ ID NO: 5) capable of directing RNA polymerase II to initiate RNA synthesis at the appropriate transcription initiation site for a particular coding sequence. A promoter may additionally comprise other recognition sequences generally positioned upstream or 5' to the TATA box, referred to as upstream promoter elements, which influence the transcription initiation rate. Such elements include a W-box (nucleotide sequence 322-327 of SEQ ID NO: 5; nucleotide sequence 364-368 and 371-375 of SEQ ID NO: 10) and a G-box (nucleotide sequence 722-727 of SEQ ID NO: 5; nucleotide sequence 415-420 of

SEQ ID NO:10). The promoter sequences of the present invention "regulate" (i.e., repress or activate) transcription from the promoter region. The regulation of transcription by the promoter sequences of the present invention is defined herein as "inducible." By "inducible" is intended the ability of the promoter sequences to  
5 regulate expression of an operably linked nucleotide sequence in response to a stimulus.

It is recognized that having identified the nucleotide sequences for the promoter regions disclosed herein, it is within the state of the art to isolate and identify additional regulatory element in the 5' untranslated region upstream from the  
10 particular promoter regions defined herein. Thus for example, the promoter regions disclosed herein may further comprise upstream regulatory elements that confer tissue-preferred expression of heterologous nucleotide sequences operably linked to the disclosed promoter sequence. See particularly, Australian Paten No. AU-A-77751/94 and U.S. Patent Nos. 5,466,785 and 5,635,618.

15 Fragments and variants of the disclosed LOX and SCIP-1 promoter nucleotide sequences are also encompassed by the present invention. By "fragment" is intended a portion of the nucleotide sequence. Fragments of a nucleotide sequence may retain biological activity and hence retain their transcriptional regulatory activity. Thus, for example, less than the entire promoter sequence disclosed herein may be utilized to  
20 drive expression of an operably linked nucleotide sequence of interest, such as a nucleotide sequence encoding a heterologous protein. Alternatively, a fragment of promoter sequence may retain the ability to regulate transcription in the presence of a stimulus when operably linked to a heterologous transcriptional initiation region. Alternatively, fragments of a nucleotide sequence that are useful as hybridization  
25 probes generally do not retain biological activity. Thus, fragments of a nucleotide sequence may range from at least about 20 nucleotides, about 50 nucleotides, about 100 nucleotides, and up to the full-length nucleotide sequence of the invention.

Thus, a fragment of a LOX or SCIP-1 promoter nucleotide sequence may encode a biologically active portion of the LOX or SCIP-1 promoter, or it may be a  
30 fragment that can be used as a hybridization probe or PCR primer using methods disclosed below. A biologically active portion of a LOX or SCIP-1 promoter can be prepared by isolating a portion of one of the LOX or SCIP-1 promoter nucleotide sequences of the invention, and assessing the activity of the portion of the LOX or

SCIP-1 promoter. Nucleic acid molecules that are fragments of a LOX or SCIP-1 promoter nucleotide sequence comprise at least 16, 20, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, or 800 nucleotides, or up to the number of nucleotides present in a full-length LOX or SCIP-1 promoter nucleotide sequence disclosed herein (for example, 880 nucleotide for SEQ ID NO: 5; 510 nucleotide for SEQ ID NO: 10). Assays to determine the activity of a promoter sequence are well known in the art. For example, a LOX or SCIP-1 promoter fragment or variant may be operably linked to the nucleotide sequence encoding any reporter protein, such as the  $\beta$ -glucuronidase protein (GUS reporter) or the luciferase protein. The DNA construct is inserted into the genome of a plant or plant cell and the mRNA or protein level of the reporter sequence is determined. See, for example, Eulgem *et al.* (1999) *EMBO. 18*: 4689-4699.

Thus, isolated sequences that have promoter activity and which hybridize under stringent conditions to the LOX and SCIP-1 sequences disclosed herein, or to fragments thereof, are encompassed by the present invention. Such sequences will be at least about 40% to 50% homologous, about 60%, 65%, or 70% homologous, and even at least about 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homologous with the disclosed sequences. That is, the sequence identity of sequences may range, sharing at least about 40% to 50%, about 60%, 65%, or 70%, and even at least about 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity.

By "variants" of the promoter sequences is intended substantially similar sequences. For nucleotide sequences naturally occurring variants such as these can be identified with the use of well-known molecular biology techniques, as, for example, with polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant nucleotide sequences also include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis. Generally, variants of a particular nucleotide sequence of the invention will have at least about 40%, 50%, 60%, 65%, 70%, generally at least about 75%, 80%, 85%, preferably at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, and more preferably at least about 98%, 99% or more sequence identity to that particular nucleotide sequence as determined by sequence alignment programs described elsewhere herein using default parameters.

The nucleotide sequences of the invention can be used to isolate corresponding sequences from other organisms, particularly other plants, more particularly other monocots. In this manner, methods such as PCR, hybridization, and the like can be used to identify such sequences based on their sequence homology to the sequences set forth herein. Sequences isolated based on their sequence identity to the entire disease resistant sequences set forth herein or to fragments thereof are encompassed by the present invention. Such sequences include sequences that are orthologs of the disclosed sequences. By "orthologs" is intended genes derived from a common ancestral gene and which are found in different species as a result of speciation. Genes found in different species are considered orthologs when their nucleotide sequences and/or their encoded protein sequences share substantial identity as defined elsewhere herein. Functions of orthologs are often highly conserved among species.

In a PCR approach, oligonucleotide primers can be designed for use in PCR reactions to amplify corresponding DNA sequences from cDNA or genomic DNA extracted from any plant of interest. Methods for designing PCR primers and PCR cloning are generally known in the art and are disclosed in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York). See also Innis *et al.*, eds. (1990) *PCR Protocols: A Guide to Methods and Applications* (Academic Press, New York); Innis and Gelfand, eds. (1995) *PCR Strategies* (Academic Press, New York); and Innis and Gelfand, eds. (1999) *PCR Methods Manual* (Academic Press, New York). Known methods of PCR include, but are not limited to, methods using paired primers, nested primers, single specific primers, degenerate primers, gene-specific primers, vector-specific primers, partially-mismatched primers, and the like.

In hybridization techniques, all or part of a known nucleotide sequence is used as a probe that selectively hybridizes to other corresponding nucleotide sequences present in a population of cloned genomic DNA fragments or cDNA fragments (*i.e.*, genomic or cDNA libraries) from a chosen organism. The hybridization probes may be genomic DNA fragments, cDNA fragments, RNA fragments, or other oligonucleotides, and may be labeled with a detectable group such as <sup>32</sup>P, or any other detectable marker. Thus, for example, probes for hybridization can be made by labeling synthetic oligonucleotides based on the disease resistant sequences of the

invention. Methods for preparation of probes for hybridization and for construction of cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

5 For example, an entire sequence disclosed herein, or one or more portions thereof, may be used as a probe capable of specifically hybridizing to the corresponding LOX, SCIP-1, rhoGAP, or ADH sequences and messenger RNAs. To achieve specific hybridization under a variety of conditions, such probes include sequences that are unique among disease resistant sequences and are preferably at  
10 least about 10 nucleotides in length, and most preferably at least about 20 nucleotides in length. Such probes may be used to amplify corresponding sequences from a chosen organism by PCR. This technique may be used to isolate additional coding sequences from a desired organism or as a diagnostic assay to determine the presence of coding sequences in an organism. Hybridization techniques include hybridization  
15 screening of plated DNA libraries (either plaques or colonies; see, for example, Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

Hybridization of such sequences may be carried out under stringent conditions. By "stringent conditions" or "stringent hybridization conditions" is  
20 intended conditions under which a probe will hybridize to its target sequence to a detectably greater degree than to other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can  
25 be identified (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

Typically, stringent conditions will be those in which the salt concentration is  
30 less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of

destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C, and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary moderate stringency  
5 conditions include hybridization in 40 to 45% formamide, 1.0 M NaCl, 1% SDS at 37°C, and a wash in 0.5X to 1X SSC at 55 to 60°C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1X SSC at 60 to 65°C. Duration of hybridization is generally less than about 24 hours, usually about 4 to about 12 hours.

10 Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the  $T_m$  can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284:  $T_m = 81.5^\circ\text{C} + 16.6 (\log M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$ ; where M is the molarity of monovalent cations, %GC is the  
15 percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe.  $T_m$  is reduced by about 1°C for each 1% of mismatching; thus,  $T_m$ ,  
20 hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with  $\geq 90\%$  identity are sought, the  $T_m$  can be decreased 10°C. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a  
25 hybridization and/or wash at 1, 2, 3, or 4°C lower than the thermal melting point ( $T_m$ ); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10°C lower than the thermal melting point ( $T_m$ ); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20°C lower than the thermal  
melting point ( $T_m$ ). Using the equation, hybridization and wash compositions, and  
30 desired  $T_m$ , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a  $T_m$  of less than 45°C (aqueous solution) or 32°C (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature

can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel *et al.*, eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2  
5 (Greene Publishing and Wiley-Interscience, New York). See Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

In general, sequences that have promotor activity or encode a rhoGAP, LOX, ADH or SCIP-1 polypeptide and which hybridize under stringent conditions to the  
10 rhoGAP, LOX, ADH or SCIP-1 sequences disclosed herein, or to fragments thereof, are encompassed by the present invention. Such sequences will be at least 40% to 50% homologous, about 60%, 65%, or 70% homologous, and even at least about 75% homologous, and even about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% homologous or more with the disclosed sequences. That is, the  
15 sequence identity of the sequences may range, sharing at least 40% to 50%, about 60%, 65%, or 70%, and even about 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity.

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides: (a) “reference sequence”, (b)  
20 “comparison window”, (c) “sequence identity”, (d) “percentage of sequence identity”, and (e) “substantial identity”.

(a) As used herein, “reference sequence” is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length cDNA or gene  
25 sequence, or the complete cDNA or gene sequence.

(b) As used herein, “comparison window” makes reference to a contiguous and specified segment of a polynucleotide sequence, wherein the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not comprise additions or deletions)  
30 for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length, and optionally can be 30, 40, 50, 100, or longer. Those of skill in the art understand that to avoid a high similarity to a



reference sequence due to inclusion of gaps in the polynucleotide sequence a gap penalty is typically introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well known in the art. Thus, the determination of percent identity between any two sequences can be accomplished using a mathematical algorithm. Non-limiting examples of such mathematical algorithms are the algorithm of Myers and Miller (1988) *CABIOS* 4:11-17; the local homology algorithm of Smith *et al.* (1981) *Adv. Appl. Math.* 2:482; the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453; the search-for-similarity-method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci.* 85:2444-2448; the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877.

Computer implementations of these mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include, but are not limited to: CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, California); the ALIGN program (Version 2.0) and GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 8 (available from Genetics Computer Group (GCG), 575 Science Drive, Madison, Wisconsin, USA). Alignments using these programs can be performed using the default parameters. The CLUSTAL program is well described by Higgins *et al.* (1988) *Gene* 73:237-244 (1988); Higgins *et al.* (1989) *CABIOS* 5:151-153; Corpet *et al.* (1988) *Nucleic Acids Res.* 16:10881-90; Huang *et al.* (1992) *CABIOS* 8:155-65; and Pearson *et al.* (1994) *Meth. Mol. Biol.* 24:307-331. The ALIGN program is based on the algorithm of Myers and Miller (1988) *supra*. A PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used with the ALIGN program when comparing amino acid sequences. The BLAST programs of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403 are based on the algorithm of Karlin and Altschul (1990) *supra*. BLAST nucleotide searches can be performed with the BLASTN program, score = 100, wordlength = 12, to obtain nucleotide sequences homologous to a nucleotide sequence encoding a protein of the invention. BLAST protein searches can be performed with the BLASTX program, score = 50, wordlength = 3, to obtain amino acid sequences homologous to a protein or polypeptide of the invention. To obtain gapped alignments for comparison

purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. See Altschul *et al.* (1997) *supra*. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. See  
5 <http://www.ncbi.nlm.nih.gov>. Alignment may also be performed manually by inspection.

Unless otherwise stated, sequence identity/similarity values provided herein  
10 refer to the value obtained using GAP Version 10 using the following parameters: % identity using GAP Weight of 50 and Length Weight of 3; % similarity using Gap Weight of 12 and Length Weight of 4, or any equivalent program. By "equivalent program" is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue  
15 matches and an identical percent sequence identity when compared to the corresponding alignment generated by the preferred program.

GAP uses the algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48: 443-453, to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. GAP considers all possible alignments  
20 and gap positions and creates the alignment with the largest number of matched bases and the fewest gaps. It allows for the provision of a gap creation penalty and a gap extension penalty in units of matched bases. GAP must make a profit of gap creation penalty number of matches for each gap it inserts. If a gap extension penalty greater than zero is chosen, GAP must, in addition, make a profit for each gap inserted of the  
25 length of the gap times the gap extension penalty. Default gap creation penalty values and gap extension penalty values in Version 10 of the Wisconsin Genetics Software Package for protein sequences are 8 and 2, respectively. For nucleotide sequences the default gap creation penalty is 50 while the default gap extension penalty is 3. The gap creation and gap extension penalties can be expressed as an integer selected  
30 from the group of integers consisting of from 0 to 200. Thus, for example, the gap creation and gap extension penalties can be 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65 or greater.

GAP presents one member of the family of best alignments. There may be many members of this family, but no other member has a better quality. GAP displays four figures of merit for alignments: Quality, Ratio, Identity, and Similarity. The Quality is the metric maximized in order to align the sequences. Ratio is the  
5 quality divided by the number of bases in the shorter segment. Percent Identity is the percent of the symbols that actually match. Percent Similarity is the percent of the symbols that are similar. Symbols that are across from gaps are ignored. A similarity is scored when the scoring matrix value for a pair of symbols is greater than or equal to 0.50, the similarity threshold. The scoring matrix used in Version 10 of the  
10 Wisconsin Genetics Software Package is BLOSUM62 (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915).

For purposes of the present invention, comparison of nucleotide or protein sequences for determination of percent sequence identity to the rhoGAP, LOX, ADH, and SCIP-1 sequences disclosed herein is preferably made using the ClustalW  
15 program (Version 1.7 or later) with its default parameters or any equivalent program. By "equivalent program" is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by the preferred program.

20 (c) As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical  
25 often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the  
30 substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity". Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage

sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE

5 (Intelligenetics, Mountain View, California).

(d) As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference  
10 sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and  
15 multiplying the result by 100 to yield the percentage of sequence identity.

(e)(i) The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70% sequence identity, preferably at least 80%, more preferably at least 90%, and most preferably at least 95%, compared to a reference sequence using one of the alignment programs  
20 described using standard parameters. One of skill in the art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning, and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least  
25 60%, more preferably at least 70%, 80%, 90%, and most preferably at least 95%.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. However, stringent  
30 conditions encompass temperatures in the range of about 1°C to about 20°C, depending upon the desired degree of stringency as otherwise qualified herein. Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides they encode are substantially identical. This

may occur, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. One indication that two nucleic acid sequences are substantially identical is when the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid.

- (e)(ii) The term "substantial identity" in the context of a peptide indicates that a peptide comprises a sequence with at least 70% sequence identity to a reference sequence, preferably 80%, more preferably 85%, most preferably at least 90% or 95% sequence identity to the reference sequence over a specified comparison window.
- Preferably, optimal alignment is conducted using the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453. An indication that two peptide sequences are substantially identical is that one peptide is immunologically reactive with antibodies raised against the second peptide. Thus, a peptide is substantially identical to a second peptide, for example, where the two peptides differ only by a conservative substitution. Peptides that are "substantially similar" share sequences as noted above except that residue positions that are not identical may differ by conservative amino acid changes.

#### Disease and Pests

- Compositions and methods for controlling pathogenic agents are provided. The anti-pathogenic compositions comprise sunflower nucleotide and polypeptide sequences. Particularly, the sunflower nucleic acid and amino acid sequences are selected from rhoGAP-1, LOX, ADH, and SCIP-1. Accordingly, the compositions and methods are useful in protecting plants against fungal pathogens, viruses, nematodes, insects and the like.

- By "disease resistance" or "pathogen resistance" is intended that the plants avoid the disease symptoms which are the outcome of plant-pathogen interactions. That is, pathogens are prevented from causing plant diseases and the associated disease symptoms, or alternatively, the disease symptoms caused by the pathogen is minimized or lessened. The methods of the invention can be utilized to protect plants from disease, particularly those diseases that are caused by plant pathogens.

By "antipathogenic compositions" is intended that the compositions of the invention have antipathogenic activity and thus are capable of suppressing,

controlling, and/or killing the invading pathogenic organism. An antipathogenic composition of the invention will reduce the disease symptoms resulting from pathogen challenge by at least about 5% to about 50%, at least about 10% to about 60%, at least about 30% to about 70%, at least about 40% to about 80%, or at least  
5 about 50% to about 90% or greater. Hence, the methods of the invention can be utilized to protect plants from disease, particularly those diseases that are caused by plant pathogens.

Assays that measure antipathogenic activity are commonly known in the art, as are methods to quantitate disease resistance in plants following pathogen infection.  
10 See, for example, U.S. Patent No. 5,614,395, herein incorporated by reference. Such techniques include, measuring over time, the average lesion diameter, the pathogen biomass, and the overall percentage of decayed plant tissues. For example, a plant either expressing an antipathogenic polypeptide or having an antipathogenic composition applied to its surface shows a decrease in tissue necrosis (*i.e.*, lesion  
15 diameter) or a decrease in plant death following pathogen challenge when compared to a control plant that was not exposed to the antipathogenic composition. Alternatively, antipathogenic activity can be measured by a decrease in pathogen biomass. For example, a plant expressing an antipathogenic polypeptide or exposed to an antipathogenic composition is challenged with a pathogen of interest. Over  
20 time, tissue samples from the pathogen-inoculated tissues are obtained and RNA is extracted. The percent of a specific pathogen RNA transcript relative to the level of a plant specific transcript allows the level of pathogen biomass to be determined. See, for example, Thomma *et al.* (1998) *Plant Biology* 95:15107-15111, herein incorporated by reference.

25 Furthermore, *in vitro* antipathogenic assays include, for example, the addition of varying concentrations of the antipathogenic composition to paper disks and placing the disks on agar containing a suspension of the pathogen of interest. Following incubation, clear inhibition zones develop around the discs that contain an effective concentration of the antipathogenic polypeptide (Liu *et al.* (1994) *Plant*  
30 *Biology* 91:1888-1892, herein incorporated by reference). Additionally, microspectrophotometrical analysis can be used to measure the *in vitro* antipathogenic properties of a composition (Hu *et al.* (1997) *Plant Mol. Biol.* 34:949-959 and

Cammue *et al.* (1992) *J. Biol. Chem.* 267: 2228-2233, both of which are herein incorporated by reference).

Methods for increasing pathogen resistance in a plant are provided. The methods involve stably transforming a plant with a DNA construct comprising an anti-pathogenic nucleotide sequence of the invention operably linked to promoter that drives expression in a plant. Such methods may find use in agriculture particularly in limiting the impact of plant pathogens on crop plants. The anti-pathogenic nucleotide sequences comprise the sunflower rhoGAP, LOX, ADH, or SCIP-1 nucleic acid molecules. While the choice of promoter will depend on the desired timing and location of expression of the anti-pathogenic nucleotide sequences, preferred promoters include constitutive and pathogen-inducible promoters.

Additionally, the compositions can be used in formulation use for their disease resistance activities. The proteins of the invention can be formulated with an acceptable carrier into a pesticidal composition(s) that is for example, a suspension, a solution, an emulsion, a dusting powder, a dispersible granule, a wettable powder, and an emulsifiable concentrate, an aerosol, an impregnated granule, an adjuvant, a coatable paste, and also encapsulations in, for example, polymer substances.

Additionally, transformed plants, plant cells, plant tissues and seeds thereof are provided.

It is understood in the art that plant DNA viruses and fungal pathogens remodel the control of the host replication and gene expression machinery to accomplish their own replication and effective infection. The present invention may be useful in preventing such corruption of the cell.

As discussed above, the sequences encoding the sunflower rhoGAP, LOX, ADH, and SCIP-1 are involved in many basic biochemical pathways and cellular functions that influence the plant defense response. Hence, the sequences of the invention may find use in disrupting cellular function of plant pathogens or insect pests as well as altering the defense mechanisms of a host plant to enhance resistance to disease or insect pests. While the invention is not bound by any particular mechanism of action, the gene products, probably proteins or polypeptides, function to inhibit or prevent plant diseases in a plant. Such gene products may be anti-pathogenic. It is recognized that the present invention is not dependent upon a particular mechanism of defense. Rather, the genes and methods of the invention

work to increase resistance of the plant to pathogens independent of how that resistance is increased or achieved.

The methods of the invention can be used with other methods available in the art for enhancing disease resistance in plants. Similarly, the plant defense mechanisms described herein may be used alone or in combination with other proteins or agents to protect against plant diseases and pathogens. Although any one of a variety of second nucleotide sequences may be utilized, specific embodiments of the invention encompass those second nucleotide sequences that, when expressed in a plant, help to increase the resistance of a plant to pathogens. It is recognized that such second nucleotide sequences may be used in either the sense or antisense orientation depending on the desired outcome. Other plant defense proteins include those described in WO 99/43823 and WO 99/43821, all of which are herein incorporated by reference.

Additionally, the LOX and SCIP-1 promoter nucleotide sequences disclosed herein are also useful for genetic engineering of plants to express a phenotype of interest. The promoter sequences may be used to regulate expression of any heterologous nucleotide sequence. Alternatively, the LOX or SCIP-1 promoter sequence may be used to drive expression of its native, i.e., naturally occurring, LOX or SCIP-1 gene sequence disclosed herein. In a specific embodiment, the LOX or SCIP-1 promoter sequences are operably linked to an anti-pathogenic nucleotide sequence and drive expression of said sequence in a plant cell. The LOX or SCIP-1 promoter sequences may therefore be used in creating or enhancing pathogen or disease resistance in a transformed plant.

Pathogens of the invention include, but are not limited to, viruses or viroids, bacteria, insects, nematodes, fungi, and the like. Viruses include any plant virus, for example, tobacco or cucumber mosaic virus, ringspot virus, necrosis virus, maize dwarf mosaic virus, etc. Specific fungal and viral pathogens for the major crops include: Soybeans: *Phytophthora megasperma* f.sp. *glycinea*, *Macrophomina phaseolina*, *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Fusarium oxysporum*, *Diaporthe phaseolorum* var. *sojae* (*Phomopsis sojae*), *Diaporthe phaseolorum* var. *caulivora*, *Sclerotium rolfsii*, *Cercospora kikuchii*, *Cercospora sojae*, *Peronospora manshurica*, *Colletotrichum dematium* (*Colletotrichum truncatum*), *Corynespora cassiicola*, *Septoria glycines*, *Phyllosticta sojicola*, *Alternaria alternata*,



- Pseudomonas syringae* p.v. *glycinea*, *Xanthomonas campestris* p.v. *phaseoli*, *Microsphaera diffusa*, *Fusarium semitectum*, *Phialophora gregata*, Soybean mosaic virus, *Glomerella glycines*, Tobacco Ring spot virus, Tobacco Streak virus, *Phakopsora pachyrhizi*, *Pythium aphanidermatum*, *Pythium ultimum*, *Pythium debaryanum*, Tomato spotted wilt virus, *Heterodera glycines* *Fusarium solani*;
- 5 Canola: *Albugo candida*, *Alternaria brassicae*, *Leptosphaeria maculans*, *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Mycosphaerella brassicicola*, *Pythium ultimum*, *Peronospora parasitica*, *Fusarium roseum*, *Alternaria alternata*; Alfalfa: *Clavibacter michiganense* subsp. *insidiosum*, *Pythium ultimum*, *Pythium irregulare*, *Pythium splendens*, *Pythium debaryanum*, *Pythium aphanidermatum*, *Phytophthora megasperma*, *Peronospora trifoliorum*, *Phoma medicaginis* var. *medicaginis*, *Cercospora medicaginis*, *Pseudopeziza medicaginis*, *Leptotrochila medicaginis*, *Fusarium*, *Xanthomonas campestris* p.v. *alfalfae*, *Aphanomyces euteiches*, *Stemphylium herbarum*, *Stemphylium alfalfae*; Wheat: *Pseudomonas syringae* p.v.
- 15 *atrofaciens*, *Urocystis agropyri*, *Xanthomonas campestris* p.v. *translucens*, *Pseudomonas syringae* p.v. *syringae*, *Alternaria alternata*, *Cladosporium herbarum*, *Fusarium graminearum*, *Fusarium avenaceum*, *Fusarium culmorum*, *Ustilago tritici*, *Ascochyta tritici*, *Cephalosporium gramineum*, *Collotetrachum graminicola*, *Erysiphe graminis* f.sp. *tritici*, *Puccinia graminis* f.sp. *tritici*, *Puccinia recondita* f.sp. *tritici*,
- 20 *Puccinia striiformis*, *Pyrenophora tritici-repentis*, *Septoria nodorum*, *Septoria tritici*, *Septoria avenae*, *Pseudocercospora herpotrichoides*, *Rhizoctonia solani*, *Rhizoctonia cerealis*, *Gaeumannomyces graminis* var. *tritici*, *Pythium aphanidermatum*, *Pythium arrhenomanes*, *Pythium ultimum*, *Bipolaris sorokiniana*, Barley Yellow Dwarf Virus, Brome Mosaic Virus, Soil Borne Wheat Mosaic Virus,
- 25 Wheat Streak Mosaic Virus, Wheat Spindle Streak Virus, American Wheat Striate Virus, *Claviceps purpurea*, *Tilletia tritici*, *Tilletia laevis*, *Ustilago tritici*, *Tilletia indica*, *Rhizoctonia solani*, *Pythium arrhenomannes*, *Pythium graminicola*, *Pythium aphanidermatum*, High Plains Virus, European wheat striate virus; Sunflower: *Orobanche*, *Plasmophora halstedii*, *Sclerotinia sclerotiorum*, Aster Yellows, *Septoria helianthi*, *Phomopsis helianthi*, *Alternaria helianthi*, *Alternaria zinniae*, *Botrytis cinerea*, *Phoma macdonaldii*, *Macrophomina phaseolina*, *Erysiphe cichoracearum*, *Rhizopus oryzae*, *Rhizopus arrhizus*, *Rhizopus stolonifer*, *Puccinia helianthi*, *Verticillium dahliae*, *Erwinia carotovorum* pv. *carotovora*, *Cephalosporium*

- acremonium*, *Phytophthora cryptogea*, *Albugo tragopogonis*; Corn: *Fusarium moniliforme* var. *subglutinans*, *Erwinia stewartii*, *Fusarium moniliforme*, *Gibberella zeae* (*Fusarium graminearum*), *Stenocarpella maydi* (*Diplodia maydis*), *Pythium irregulare*, *Pythium debaryanum*, *Pythium graminicola*, *Pythium splendens*, *Pythium ultimum*, *Pythium aphanidermatum*, *Aspergillus flavus*, *Bipolaris maydis* O, T (*Cochliobolus heterostrophus*), *Helminthosporium carbonum* I, II & III (*Cochliobolus carbonum*), *Exserohilum turcicum* I, II & III, *Helminthosporium pedicellatum*, *Physoderma maydis*, *Phyllosticta maydis*, *Kabatiella-maydis*, *Cercospora sorghi*, *Ustilago maydis*, *Puccinia sorghi*, *Puccinia polysora*, *Macrophomina phaseolina*,  
 5 *Penicillium oxalicum*, *Nigrospora oryzae*, *Cladosporium herbarum*, *Curvularia lunata*, *Curvularia inaequalis*, *Curvularia pallescens*, *Clavibacter michiganense* subsp. *nebraskense*, *Trichoderma viride*, Maize Dwarf Mosaic Virus A & B, Wheat Streak Mosaic Virus, Maize Chlorotic Dwarf Virus, *Claviceps sorghi*, *Pseudomonas avenae*, *Erwinia chrysanthemi* pv. *zea*, *Erwinia carotovora*, Corn stunt Spiroplasma,  
 10 *Diplodia macrospora*, *Sclerophthora macrospora*, *Peronosclerospora sorghi*, *Peronosclerospora philippinensis*, *Peronosclerospora maydis*, *Peronosclerospora sacchari*, *Sphacelotheca reiliana*, *Physopella zeae*, *Cephalosporium maydis*, *Cephalosporium acremonium*, Maize Chlorotic Mottle Virus, High Plains Virus, Maize Mosaic Virus, Maize Rayado Fino Virus, Maize Streak Virus, Maize Stripe  
 20 Virus, Maize Rough Dwarf Virus; Sorghum: *Exserohilum turcicum*, *Colletotrichum graminicola* (*Glomerella graminicola*), *Cercospora sorghi*, *Gloeocercospora sorghi*, *Ascochyta sorghina*, *Pseudomonas syringae* p.v. *syringae*, *Xanthomonas campestris* p.v. *holcicola*, *Pseudomonas andropogonis*, *Puccinia purpurea*, *Macrophomina phaseolina*, *Perconia circinata*, *Fusarium moniliforme*, *Alternaria alternata*,  
 25 *Bipolaris sorghicola*, *Helminthosporium sorghicola*, *Curvularia lunata*, *Phoma insidiosa*, *Pseudomonas avenae* (*Pseudomonas alboprecipitans*), *Ramulispora sorghi*, *Ramulispora sorghicola*, *Phyllachara sacchari*, *Sporisorium reilianum* (*Sphacelotheca reiliana*), *Sphacelotheca cruenta*, *Sporisorium sorghi*, Sugarcane mosaic H, Maize Dwarf Mosaic Virus A & B, *Claviceps sorghi*, *Rhizoctonia solani*,  
 30 *Acremonium strictum*, *Sclerophthona macrospora*, *Peronosclerospora sorghi*, *Peronosclerospora philippinensis*, *Sclerospora graminicola*, *Fusarium graminearum*, *Fusarium oxysporum*, *Pythium arrhenomanes*, *Pythium graminicola*, etc.,

Nematodes include parasitic nematodes such as root-knot, cyst, lesion, and renniform nematodes, etc.

- Insect pests include insects selected from the orders Coleoptera, Diptera, Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera,
- 5 Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, etc., particularly Coleoptera and Lepidoptera. Insect pests of the invention for the major crops include: Maize: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Helicoverpa zea*, corn earworm; *Spodoptera frugiperda*, fall armyworm; *Diatraea grandiosella*, southwestern corn borer; *Elasmopalpus lignosellus*, lesser
- 10 cornstalk borer; *Diatraea saccharalis*, sugarcane borer; *Diabrotica virgifera*, western corn rootworm; *Diabrotica longicornis barberi*, northern corn rootworm; *Diabrotica undecimpunctata howardi*, southern corn rootworm; *Melanotus spp.*, wireworms; *Cyclocephala borealis*, northern masked chafer (white grub); *Cyclocephala immaculata*, southern masked chafer (white grub); *Popillia japonica*, Japanese beetle;
- 15 *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Anuraphis maidiradicis*, corn root aphid; *Blissus leucopterus leucopterus*, chinch bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Hylemya platura*, seedcorn maggot; *Agromyza parvicornis*, corn blot leafminer; *Anaphothrips*
- 20 *obscurus*, grass thrips; *Solenopsis milesta*, thief ant; *Tetranychus urticae*, twospotted spider mite; Sorghum: *Chilo partellus*, sorghum borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Feltia subterranea*, granulate cutworm; *Phyllophaga crinita*, white grub; *Eleodes*, *Conoderus*, and *Aeolus spp.*, wireworms; *Oulema melanopus*, cereal
- 25 leaf beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Sipha flava*, yellow sugarcane aphid; *Blissus leucopterus leucopterus*, chinch bug; *Contarinia sorghicola*, sorghum midge; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Wheat: *Pseudaletia unipunctata*, army worm; *Spodoptera frugiperda*,
- 30 fall armyworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Agrotis orthogonia*, western cutworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Oulema melanopus*, cereal leaf beetle; *Hypera punctata*, clover leaf weevil; *Diabrotica undecimpunctata howardi*, southern corn rootworm; Russian wheat aphid; *Schizaphis*

- graminum*, greenbug; *Macrosiphum avenae*, English grain aphid; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Mayetiola destructor*, Hessian fly; *Sitodiplosis mosellana*, wheat midge; *Meromyza americana*, wheat stem
- 5 maggot; *Hylemya coarctata*, wheat bulb fly; *Frankliniella fusca*, tobacco thrips; *Cephus cinctus*, wheat stem sawfly; *Aceria tulipae*, wheat curl mite; Sunflower: *Suleima helianthana*, sunflower bud moth; *Homoeosoma electellum*, sunflower moth; *zygogramma exclamationis*, sunflower beetle; *Bothyrus gibbosus*, carrot beetle; *Neolasioptera murtfeldtiana*, sunflower seed midge; Cotton: *Heliothis virescens*,
- 10 cotton budworm; *Helicoverpa zea*, cotton bollworm; *Spodoptera exigua*, beet armyworm; *Pectinophora gossypiella*, pink bollworm; *Anthonomus grandis grandis*, boll weevil; *Aphis gossypii*, cotton aphid; *Pseudatomoscelis seriatus*, cotton fleahopper; *Trialeurodes abutilonea*, bandedwinged whitefly; *Lygus lineolaris*, tarnished plant bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus*
- 15 *differentialis*, differential grasshopper; *Thrips tabaci*, onion thrips; *Frankliniella fusca*, tobacco thrips; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Rice: *Diatraea saccharalis*, sugarcane borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Colaspis brunnea*, grape colaspis; *Lissorhoptrus oryzophilus*, rice water weevil; *Sitophilus*
- 20 *oryzae*, rice weevil; *Nephotettix nigropictus*, rice leafhopper; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug; Soybean: *Pseudoplusia includens*, soybean looper; *Anticarsia gemmatilis*, velvetbean caterpillar; *Plathypena scabra*, green cloverworm; *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Spodoptera exigua*, beet armyworm; *Heliothis*
- 25 *virescens*, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Epilachna varivestis*, Mexican bean beetle; *Myzus persicae*, green peach aphid; *Empoasca fabae*, potato leafhopper; *Acrosternum hilare*, green stink bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Hylemya platura*, seedcorn maggot; *Sericothrips variabilis*, soybean thrips; *Thrips tabaci*, onion thrips;
- 30 *Tetranychus turkestanii*, strawberry spider mite; *Tetranychus urticae*, twospotted spider mite; Barley: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Schizaphis graminum*, greenbug; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug; *Euschistus servus*, brown stink bug; *Delia*

*platura*, seedcorn maggot; *Mayetiola destructor*, Hessian fly; *Petrobia latens*, brown wheat mite; Oil Seed Rape: *Brevicoryne brassicae*, cabbage aphid; *Phyllotreta cruciferae*, Flea beetle; *Mamestra configurata*, Bertha armyworm; *Plutella xylostella*, Diamond-back moth; *Delia* ssp., Root maggots.

5

### Expression of Sequences

The nucleic acid sequences of the present invention can be expressed in a host cell such as bacteria, yeast, insect, mammalian, or preferably plant cells. It is expected that those of skill in the art are knowledgeable in the numerous expression systems available for expression of a nucleic acid encoding a protein of the present invention. No attempt to describe in detail the various methods known for the expression of proteins in prokaryotes or eukaryotes will be made.

As used herein, "heterologous" in reference to a nucleic acid is a nucleic acid that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. For example, a promoter operably linked to a heterologous structural gene is from a species different from that from which the structural gene was derived, or, if from the same species, one or both are substantially modified from their original form. A heterologous protein may originate from a foreign species, or, if from the same species, is substantially modified from its original form by deliberate human intervention.

By "host cell" is meant a cell, which comprises a heterologous nucleic acid sequence of the invention. Host cells may be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells. Preferably, host cells are monocotyledonous or dicotyledonous plant cells. A particularly preferred monocotyledonous host cell is a maize host cell.

The disease resistant sequences of the invention are provided in expression cassettes or DNA constructs for expression in the plant of interest. The cassette will include 5' and 3' regulatory sequences operably linked to a rhoGAP, LOX, ADH, or SCIP-1 sequence of the invention. By "operably linked" is intended a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being

linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame. The cassette may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes.

- 5       Such an expression cassette is provided with a plurality of restriction sites for insertion of the disease resistant sequence to be under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes.

- 10       The expression cassette will include in the 5'-3' direction of transcription, a transcriptional and translational initiation region, a disease resistant DNA sequence of the invention, and a transcriptional and translational termination region functional in plants. The transcriptional initiation region, the promoter, may be native or analogous or foreign or heterologous to the plant host. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. By "foreign" is intended that  
15       the transcriptional initiation region is not found in the native plant into which the transcriptional initiation region is introduced. As used herein, a chimeric gene comprises a coding sequence operably linked to a transcription initiation region that is heterologous to the coding sequence.

- 20       While it may be preferable to express the sequences using heterologous promoters, the native promoter sequences may be used. Such constructs would change expression levels of the disease resistant RNA/protein in the plant or plant cell. Thus, the phenotype of the plant or plant cell is altered.

- 25       The termination region may be native with the transcriptional initiation region, may be native with the operably linked DNA sequence of interest, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also Guerineau *et al.* (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell* 2:1261-1272; Munroe *et al.* (1990) *Gene* 91:151-158;  
30       Ballas *et al.* (1989) *Nucleic Acids Res.* 17:7891-7903; and Joshi *et al.* (1987) *Nucleic Acid Res.* 15:9627-9639.

Where appropriate, the gene(s) may be optimized for increased expression in the transformed plant. That is, the genes can be synthesized using plant-preferred

codons for improved expression. Methods are available in the art for synthesizing plant-preferred genes. See, for example, U.S. Patent Nos. 5,380,831, and 5,436,391, and Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

- 5 Additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats, and other such well-characterized sequences that may be deleterious to gene expression. The G-C content of the sequence may be adjusted to levels average for a given
- 10 cellular host, as calculated by reference to known genes expressed in the host cell. When possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures.

- The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation.
- 15 Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein *et al.* (1989) *PNAS USA* 86:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Allison *et al.* (1986); MDMV leader (Maize Dwarf Mosaic Virus); *Virology* 154:9-20), and human immunoglobulin heavy-chain binding protein
- 20 (BiP), (Macejak *et al.* (1991) *Nature* 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4) (Jobling *et al.* (1987) *Nature* 325:622-625); tobacco mosaic virus leader (TMV) (Gallie *et al.* (1989) in *Molecular Biology of RNA*, ed. Cech (Liss, New York), pp. 237-256); and maize chlorotic mottle virus leader (MCMV) (Lommel *et al.* (1991) *Virology* 81:382-385). See also,
- 25 Della-Cioppa *et al.* (1987) *Plant Physiol.* 84:965-968. Other methods known to enhance translation can also be utilized, for example, introns, and the like.

- In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be
- 30 employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in vitro* mutagenesis, primer repair,

restriction, annealing, resubstitutions, e.g., transitions and transversions, may be involved.

Generally, the expression cassette will comprise a selectable marker gene for the selection of transformed cells. Selectable marker genes are utilized for the selection of transformed cells or tissues. Marker genes include genes encoding antibiotic resistance, such as those encoding neomycin phosphotransferase II (NEO) and hygromycin phosphotransferase (HPT), as well as genes conferring resistance to herbicidal compounds, such as glufosinate ammonium, bromoxynil, imidazolinones, and 2,4-dichlorophenoxyacetate (2,4-D). See generally, Yarranton (1992) *Curr. Opin. Biotech.* 3:506-511; Christopherson *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:6314-6318; Yao *et al.* (1992) *Cell* 71:63-72; Reznikoff (1992) *Mol. Microbiol.* 6:2419-2422; Barkley *et al.* (1980) in *The Operon*, pp. 177-220; Hu *et al.* (1987) *Cell* 48:555-566; Brown *et al.* (1987) *Cell* 49:603-612; Figge *et al.* (1988) *Cell* 52:713-722; Deuschle *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:5400-5404; Fuerst *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:2549-2553; Deuschle *et al.* (1990) *Science* 248:480-483; Gossen (1993) Ph.D. Thesis, University of Heidelberg; Reines *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:1917-1921; Labow *et al.* (1990) *Mol. Cell. Biol.* 10:3343-3356; Zambretti *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:3952-3956; Baim *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:5072-5076; Wyborski *et al.* (1991) *Nucleic Acids Res.* 19:4647-4653; Hillenand-Wissman (1989) *Topics Mol. Struc. Biol.* 10:143-162; Degenkolb *et al.* (1991) *Antimicrob. Agents Chemother.* 35:1591-1595; Kleinschmidt *et al.* (1988) *Biochemistry* 27:1094-1104; Bonin (1993) Ph.D. Thesis, University of Heidelberg; Gossen *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Oliva *et al.* (1992) *Antimicrob. Agents Chemother.* 36:913-919; Hlavka *et al.* (1985) *Handbook of Experimental Pharmacology*, Vol. 78 (Springer-Verlag, Berlin); Gill *et al.* (1988) *Nature* 334:721-724. Such disclosures are herein incorporated by reference.

The above list of selectable marker genes is not meant to be limiting. Any selectable marker gene can be used in the present invention.

A number of promoters can be used in the practice of the invention. The promoters can be selected based on the desired outcome. That is, the nucleic acids can be combined with constitutive, tissue-preferred, or other promoters for expression in plants. Such constitutive promoters include, for example, the core promoter of the Rsyn7 (PCT Application Serial No. US99/03863); Scp1 promoter (U.S. Patent



Application Serial No. 09/028,819), rice actin (McElroy *et al.* (1990) *Plant Cell* 2:163-171); ubiquitin (Christensen *et al.* (1989) *Plant Mol. Biol.* 12:619-632 and Christensen *et al.* (1992) *Plant Mol. Biol.* 18:675-689); pEMU (Last *et al.* (1991) *Theor. Appl. Genet.* 81:581-588); MAS (Velten *et al.* (1984) *EMBO J.* 3:2723-2730);  
5 ALS promoter (U.S. Application Serial No. 08/409,297), and the like. Other constitutive promoters include, for example, U.S. Patent Nos. 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; and 5,608,142.

Generally, it will be beneficial to express the gene from an inducible promoter, particularly from a pathogen-inducible promoter. Such promoters include those from  
10 pathogenesis-related proteins (PR proteins), which are induced following infection by a pathogen; e.g., PR proteins, SAR proteins, beta-1,3-glucanase, chitinase, etc. See, for example, Redolfi *et al.* (1983) *Neth. J. Plant Pathol.* 89:245-254; Uknes *et al.* (1992) *Plant Cell* 4:645-656; and Van Loon (1985) *Plant Mol. Virol.* 4:111-116. See also the copending application entitled "Inducible Maize Promoters", U.S. Patent  
15 Application Serial No. 09/257,583, filed February 25, 1999.

Of interest are promoters that are expressed locally at or near the site of pathogen infection. See, for example, Marineau *et al.* (1987) *Plant Mol. Biol.* 9:335-342; Matton *et al.* (1989) *Molecular Plant-Microbe Interactions* 2:325-331; Somsisch *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:2427-2430; Somsisch *et al.* (1988) *Mol.*  
20 *Gen. Genet.* 2:93-98; and Yang (1996) *Proc. Natl. Acad. Sci. USA* 93:14972-14977. See also, Chen *et al.* (1996) *Plant J.* 10:955-966; Zhang *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:2507-2511; Warner *et al.* (1993) *Plant J.* 3:191-201; Siebertz *et al.* (1989) *Plant Cell* 1:961-968; U.S. Patent No. 5,750,386 (nematode-inducible); and the references cited therein. Of particular interest is the inducible promoter for the  
25 maize PRms gene, whose expression is induced by the pathogen *Fusarium moniliforme* (see, for example, Cordero *et al.* (1992) *Physiol. Mol. Plant Path.* 41:189-200).

Additionally, as pathogens find entry into plants through wounds or insect damage, a wound-inducible promoter may be used in the constructions of the  
30 invention. Such wound-inducible promoters include potato proteinase inhibitor (pin II) gene (Ryan (1990) *Ann. Rev. Phytopath.* 28:425-449; Duan *et al.* (1996) *Nature Biotechnology* 14:494-498); wun1 and wun2, US Patent No. 5,428,148; win1 and win2 (Stanford *et al.* (1989) *Mol. Gen. Genet.* 215:200-208); systemin (McGurl *et al.*

(1992) *Science* 225:1570-1573); WIP1 (Rohmeier *et al.* (1993) *Plant Mol. Biol.* 22:783-792; Eckelkamp *et al.* (1993) *FEBS Letters* 323:73-76); MPI gene (Corderok *et al.* (1994) *Plant J.* 6(2):141-150); and the like, herein incorporated by reference.

5 Chemical-regulated promoters can be used to modulate the expression of a gene in a plant through the application of an exogenous chemical regulator.

Depending upon the objective, the promoter may be a chemical-inducible promoter, where application of the chemical induces gene expression, or a chemical-repressible promoter, where application of the chemical represses gene expression. Chemical-inducible promoters are known in the art and include, but are not limited to, the maize  
10 In2-2 promoter, which is activated by benzenesulfonamide herbicide safeners, the maize GST promoter, which is activated by hydrophobic electrophilic compounds that are used as pre-emergent herbicides, and the tobacco PR-1a promoter, which is activated by salicylic acid. Other chemical-regulated promoters of interest include steroid-responsive promoters (see, for example, the glucocorticoid-inducible promoter  
15 in Schena *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:10421-10425 and McNellis *et al.* (1998) *Plant J.* 14(2):247-257) and tetracycline-inducible and tetracycline-repressible promoters (see, for example, Gatz *et al.* (1991) *Mol. Gen. Genet.* 227:229-237, and U.S. Patent Nos. 5,814,618 and 5,789,156), herein incorporated by reference.

20 The method of transformation/transfection is not critical to the instant invention; various methods of transformation or transfection are currently available. Thus, any method, which provides for effective transformation/transfection may be employed. Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e.,  
25 monocot or dicot, targeted for transformation. Suitable methods of introducing nucleotide sequences into plant cells and subsequent insertion into the plant genome include microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334), electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606, *Agrobacterium*-mediated transformation (Townsend *et al.*, U.S. Patent No. 5,563,055; Zhao *et al.*, U.S. Patent No. 5,981,840), direct gene transfer (Paszkowski *et al.* (1984) *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford  
30 *et al.*, U.S. Patent No. 4,945,050; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and*

- Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe *et al.* (1988) *Biotechnology* 6:923-926). Also see Weissinger *et al.* (1988) *Ann. Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant Physiol.* 87:671-674
- 5 (soybean); McCabe *et al.* (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.* 27P:175-182 (soybean); Singh *et al.* (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *Biotechnology* 6:559-563 (maize); Tomes, U.S. Patent
- 10 No. 5,240,855; Busing *et al.*, U.S. Patent Nos. 5,322,783 and 5,324,646; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg (Springer-Verlag, Berlin) (maize); Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *Biotechnology* 8:833-839 (maize);
- 15 Hooykaas-Van Slogteren *et al.* (1984) *Nature (London)* 311:763-764; Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet *et al.* (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman *et al.* (Longman, New York), pp. 197-209 (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418 and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated
- 20 transformation); D'Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation); Li *et al.* (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.* (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference.
- 25       The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick *et al.* (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic identified. Two or
- 30 more generations may be grown to ensure that constitutive expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure constitutive expression of the desired phenotypic characteristic has been achieved.

The present invention may be used for transformation of any plant species, including, but not limited to, monocots and dicots. Examples of plants of interest include, but are not limited to, corn (*Zea mays*), *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa (*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet (e.g., pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria italica*), finger millet (*Eleusine coracana*)), sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), wheat (*Triticum aestivum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Coffea* spp.), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees (*Citrus* spp.), cocoa (*Theobroma cacao*), tea (*Camellia sinensis*), banana (*Musa* spp.), avocado (*Persea americana*), fig (*Ficus casica*), guava (*Psidium guajava*), mango (*Mangifera indica*), olive (*Olea europaea*), papaya (*Carica papaya*), cashew (*Anacardium occidentale*), macadamia (*Macadamia integrifolia*), almond (*Prunus amygdalus*), sugar beets (*Beta vulgaris*), sugarcane (*Saccharum* spp.), oats, barley, vegetables, ornamentals, and conifers.

Vegetables include tomatoes (*Lycopersicon esculentum*), lettuce (e.g., *Lactuca sativa*), green beans (*Phaseolus vulgaris*), lima beans (*Phaseolus limensis*), peas (*Lathyrus* spp.), and members of the genus *Cucumis* such as cucumber (*C. sativus*), cantaloupe (*C. cantalupensis*), and musk melon (*C. melo*). Ornamentals include azalea (*Rhododendron* spp.), hydrangea (*Macrophylla hydrangea*), hibiscus (*Hibiscus rosasanensis*), roses (*Rosa* spp.), tulips (*Tulipa* spp.), daffodils (*Narcissus* spp.), petunias (*Petunia hybrida*), carnation (*Dianthus caryophyllus*), poinsettia (*Euphorbia pulcherrima*), and chrysanthemum. Conifers that may be employed in practicing the present invention include, for example, pines such as loblolly pine (*Pinus taeda*), slash pine (*Pinus elliotii*), ponderosa pine (*Pinus ponderosa*), lodgepole pine (*Pinus contorta*), and Monterey pine (*Pinus radiata*); Douglas-fir (*Pseudotsuga menziesii*); Western hemlock (*Tsuga canadensis*); Sitka spruce (*Picea glauca*); redwood (*Sequoia sempervirens*); true firs such as silver fir (*Abies amabilis*) and balsam fir (*Abies balsamea*); and cedars such as Western red cedar (*Thuja plicata*) and Alaska yellow-cedar (*Chamaecyparis nootkatensis*). Preferably, plants of the present invention

are crop plants (for example, corn, alfalfa, sunflower, *Brassica*, soybean, cotton, safflower, peanut, sorghum, wheat, millet, tobacco, etc.), more preferably corn and soybean plants, yet more preferably corn plants.

Prokaryotic cells may be used as hosts for expression. Prokaryotes most frequently are represented by various strains of *E. coli*; however, other microbial strains may also be used. Commonly used prokaryotic control sequences which are defined herein to include promoters for transcription initiation, optionally with an operator, along with ribosome binding sequences, include such commonly used promoters as the beta lactamase (penicillinase) and lactose (lac) promoter systems (Chang *et al.* (1977) *Nature* 198:1056), the tryptophan (trp) promoter system (Goeddel *et al.* (1980) *Nucleic Acids Res.* 8:4057) and the lambda derived P L promoter and N-gene ribosome binding site (Shimatake *et al.* (1981) *Nature* 292:128). The inclusion of selection markers in DNA vectors transfected in *E. coli* is also useful. Examples of such markers include genes specifying resistance to ampicillin, tetracycline, or chloramphenicol.

The vector is selected to allow introduction into the appropriate host cell. Bacterial vectors are typically of plasmid or phage origin. Appropriate bacterial cells are infected with phage vector particles or transfected with naked phage vector DNA. If a plasmid vector is used, the bacterial cells are transfected with the plasmid vector DNA. Expression systems for expressing a protein of the present invention are available using *Bacillus sp.* and *Salmonella* (Palva *et al.* (1983) *Gene* 22:229-235); Mosbach *et al.* (1983) *Nature* 302:543-545).

A variety of eukaryotic expression systems such as yeast, insect cell lines, plant and mammalian cells, are known to those of skill in the art. As explained briefly below, a polynucleotide of the present invention can be expressed in these eukaryotic systems. In some embodiments, transformed/transfected plant cells, as discussed *infra*, are employed as expression systems for production of the proteins of the instant invention.

Synthesis of heterologous nucleotide sequences in yeast is well known (Sherman *et al.* (1982) *Methods in Yeast Genetics*, Cold Spring Harbor Laboratory). Two widely utilized yeasts for production of eukaryotic proteins are *Saccharomyces cerevisiae* and *Pichia pastoris*. Vectors, strains, and protocols for expression in *Saccharomyces* and *Pichia* are known in the art and available from commercial

suppliers (e.g., Invitrogen). Suitable vectors usually have expression control sequences, such as promoters, including 3-phosphoglycerate kinase or alcohol oxidase, and an origin of replication, termination sequences and the like as desired.

5 A protein of the present invention, once expressed, can be isolated from yeast by lysing the cells and applying standard protein isolation techniques to the lists. The monitoring of the purification process can be accomplished by using Western blot techniques or radioimmunoassay of other standard immunoassay techniques.

The sequences of the present invention can also be ligated to various expression vectors for use in transfecting cell cultures of, for instance, mammalian, 10 insect, or plant origin. Illustrative cell cultures useful for the production of the peptides are mammalian cells. A number of suitable host cell lines capable of expressing intact proteins have been developed in the art, and include the HEK293, BHK21, and CHO cell lines. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter (e.g. the 15 CMV promoter, a HSV *tk* promoter or *pgk* (phosphoglycerate kinase) promoter), an enhancer (Queen *et al.* (1986) *Immunol. Rev.* 89:49), and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites (e.g., an SV40 large T Ag poly A addition site), and transcriptional terminator sequences. Other animal cells useful for production of proteins of the present 20 invention are available, for instance, from the American Type Culture Collection.

Appropriate vectors for expressing proteins of the present invention in insect cells are usually derived from the SF9 baculovirus. Suitable insect cell lines include mosquito larvae, silkworm, armyworm, moth and *Drosophila* cell lines such as a Schneider cell line (See, Schneider (1987) *J. Embryol. Exp. Morphol.* 27:353-365).

25 As with yeast, when higher animal or plant host cells are employed, polyadenylation or transcription terminator sequences are typically incorporated into the vector. An example of a terminator sequence is the polyadenylation sequence from the bovine growth hormone gene. Sequences for accurate splicing of the transcript may also be included. An example of a splicing sequence is the VP1 intron 30 from SV40 (Sprague *et al.* (1983) *J. Virol.* 45:773-781). Additionally, gene sequences to control replication in the host cell may be incorporated into the vector such as those found in bovine papilloma virus type-vectors (Saveria-Campo (1985) *DNA Cloning*

*Vol. II a Practical Approach*, D.M. Glover, Ed., IRL Press, Arlington, Virginia, pp. 213-238).

Animal and lower eukaryotic (e.g., yeast) host cells are competent or rendered competent for transfection by various means. There are several well-known methods of introducing DNA into animal cells. These include: calcium phosphate precipitation, fusion of the recipient cells with bacterial protoplasts containing the DNA, treatment of the recipient cells with liposomes containing the DNA, DEAE dextrin, electroporation, biolistics, and micro-injection of the DNA directly into the cells. The transfected cells are cultured by means well known in the art (Kuchler (1997) *Biochemical Methods in Cell Culture and Virology*, Dowden, Hutchinson and Ross, Inc.).

It is recognized that with these nucleotide sequences, antisense constructions, complementary to at least a portion of the messenger RNA (mRNA) for the disease resistant sequences can be constructed. Antisense nucleotides are constructed to hybridize with the corresponding mRNA. Modifications of the antisense sequences may be made as long as the sequences hybridize to and interfere with expression of the corresponding mRNA. In this manner, antisense constructions having 70%, preferably 80%, more preferably 85% sequence identity to the corresponding antisensed sequences may be used. Furthermore, portions of the antisense nucleotides may be used to disrupt the expression of the target gene. Generally, sequences of at least 50 nucleotides, 100 nucleotides, 200 nucleotides, or greater may be used.

The nucleotide sequences of the present invention may also be used in the sense orientation to suppress the expression of endogenous genes in plants. Methods for suppressing gene expression in plants using nucleotide sequences in the sense orientation are known in the art. The methods generally involve transforming plants with a DNA construct comprising a promoter that drives expression in a plant operably linked to at least a portion of a nucleotide sequence that corresponds to the transcript of the endogenous gene. Typically, such a nucleotide sequence has substantial sequence identity to the sequence of the transcript of the endogenous gene, preferably greater than about 65% sequence identity, more preferably greater than about 85% sequence identity, most preferably greater than about 95% sequence identity. See, U.S. Patent Nos. 5,283,184 and 5,034,323; herein incorporated by reference.

The nucleotide sequences of the present invention may also be used in the sense orientation to suppress the expression of endogenous genes in plants. Methods for suppressing gene expression in plants using nucleotide sequences in the sense orientation are known in the art. The methods generally involve transforming plants with a DNA construct comprising a promoter that drives expression in a plant operably linked to at least a portion of a nucleotide sequence that corresponds to the transcript of the endogenous gene. Typically, such a nucleotide sequence has substantial sequence identity to the sequence of the transcript of the endogenous gene, preferably greater than about 65% sequence identity, more preferably greater than about 85% sequence identity, most preferably greater than about 95% sequence identity. See, U.S. Patent Nos. 5,283,184 and 5,034,323; herein incorporated by reference.

In some embodiments, the content and/or composition of polypeptides of the present invention in a plant may be modulated by altering, *in vivo* or *in vitro*, the promoter of a gene to up- or down- regulate gene expression. In some embodiments, the coding regions of native genes of the present invention can be altered via substitution, addition, insertion, or deletion to decrease activity of the encoded enzyme. See, e.g., Kmiec, U.S. Patent 5,565,350; Zarling *et al.*, PCT/US93/03868. And in some embodiments, an isolated nucleic acid (e.g., a vector) comprising a promoter sequence is transfected into a plant cell. Subsequently, a plant cell comprising the promoter operably linked to a polynucleotide of the present invention is selected for by means known to those of skill in the art such as, but not limited to, Southern blot, DNA sequencing, or PCR analysis using primers specific to the promoter and to the gene and detecting amplicons produced therefrom. A plant or plant part altered or modified by the foregoing embodiments is grown under plant forming conditions for a time sufficient to modulate the concentration and/or composition of polypeptides of the present invention in the plant. Plant forming conditions are well known in the art and discussed briefly, *supra*.

In general, concentration or composition is increased or decreased by at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% relative to a native control plant, plant part, or cell lacking the aforementioned recombinant expression cassette. Modulation in the present invention may occur during and/or subsequent to growth of the plant to the desired stage of development. Modulating nucleic acid expression



temporally and/or in particular tissues can be controlled by employing the appropriate promoter operably linked to a polynucleotide of the present invention in, for example, sense or antisense orientation as discussed in greater detail, *supra*. Induction of expression of a polynucleotide of the present invention can also be controlled by  
5 exogenous administration of an effective amount of inducing compound. Inducible promoters and inducing compounds, which activate expression from these promoters, are well known in the art. In preferred embodiments, the polypeptides of the present invention are modulated in monocots, particularly maize.

#### 10 Methods of Use for LOX and SCIP-1 Promoter Sequences

The nucleotide sequences for the LOX and SCIP-1 promoters disclosed in the present invention, as well as variants and fragments thereof, are useful in the genetic manipulation of any host cell, preferably plant cell, when assembled with a DNA construct such that the promoter sequence is operably linked to a nucleotide sequence  
15 encoding a heterologous protein of interest. In this manner, the nucleotide sequences of the LOX and SCIP-1 promoter of the invention are provided in expression cassettes along with heterologous nucleotide sequences for expression in the host cell of interest.

The promoters for the LOX and SCIP-1 genes may regulate expression of  
20 operably linked nucleotide sequences in an inducible manner. That is, expression of the operably linked nucleotide sequences in a host cell (i.e., plant cell) is induced in response to a stimulus. By "stimulus" is intended a chemical, which may be applied externally or may accumulate in response to another external stimulus; a pathogen, which may, for example, induce expression as a result of invading a plant cell; or  
25 other factor such as environmental stresses, including but not limited to, drought, temperature, and salinity.

As such, the stimulus either directly or indirectly regulates the activity (i.e., an increase in initiation or expression) of an inducible promoter. By "direct action" is intended that the stimulus regulates transcription via a direct interaction between the  
30 stimulus and the DNA sequence. By "indirect action" is meant that the regulation occurs via an interaction between the stimulus and some other endogenous or exogenous component in the system, the ultimate result of the indirect action being regulation of the inducible promoter. The stimulus can result from a biotic or abiotic

stress, including for example, tissue wounding (i.e., insect herbivory, wind, intentional abiotic infliction of tissue injury or wounding for the purpose of experimentation and/or expression analysis); wound-responsive chemicals (i.e., chemicals that result in the activation of wound-response signal transduction pathways, including, various hormones, jasmonic acid, abscissic acid, linolenic acid, ethylene, their chemical analogues, derivatives, precursors, and the like); pathogens (i.e., fungi, bacteria, nematodes, mycoplasmas, viruses, and insects and the like); and various environmental stresses (i.e., heat, drought, cold, reactive oxygen species and/or radiation). Hence, the promoter of the present invention can be used in combination with a nucleotide sequence that enhances disease resistance, and the compositions therefor find use in the defense of a plant against disease, pathogens, and the like.

Synthetic hybrid promoter regions are known in the art. Such regions comprise upstream promoter elements of one nucleotide sequence operably linked to the promoter element of another nucleotide sequence. In an embodiment of the invention, heterologous gene expression is controlled by a synthetic hybrid promoter comprising the LOX or SCIP-1 promoter sequences of the invention, or a variant or fragment thereof, operably linked to upstream promoter element(s) from a heterologous promoter. Upstream promoter elements that are involved in the plant defense system have been identified and may be used to generate a synthetic promoter. See, for example, Rushton *et al.* (1998) *Curr. Opin. Plant Biol.* 1:311-315. Alternatively, a synthetic LOX or SCIP-1 promoter sequence may comprise duplications of the upstream promoter elements found within the LOX or SCIP-1 promoter sequence. Such elements include, for example the G-box (nucleotides 722-727 of SEQ ID NO: 5; nucleotides 415-420 of SEQ ID NO: 10) or W-box (nucleotides 322-327 of SEQ ID NO: 5; nucleotides 364-368 and 371-375 of SEQ ID NO: 10).

It is recognized that the promoter sequence of the invention may be used with its native LOX or SCIP-1 coding sequences. A DNA construct comprising the LOX or SCIP-1 promoter operably linked with its native LOX or SCIP-1 gene may be used to transform any plant of interest to bring about a desired phenotypic change, such as enhanced disease resistance. Where the promoter and its native gene is naturally occurring within the plant, i.e., in sunflower, transformation of the plant with these

operably linked sequences also results in either a change in phenotype, such as enhanced disease resistance, the insertion of operably linked sequences within a different region of the chromosome thereby altering the plant's genome, or the modulation in the level of expression of the nucleotide sequence of interest.

5           In another embodiment of the invention, expression cassettes will comprise a transcriptional initiation region comprising the LOX or SCIP-1 promoter nucleotide sequences disclosed herein, or variants or fragments thereof, operably linked to the heterologous nucleotide sequence whose expression is to be controlled by the inducible promoter of the invention.

10           The promoter nucleotide sequences and methods disclosed herein are useful in regulating expression of any heterologous nucleotide sequence in a host plant in order to vary the phenotype of a plant. Various changes in phenotype are of interest including modifying the fatty acid composition in a plant, altering the amino acid content of a plant, altering a plant's pathogen defense mechanism, and the like. These  
15 results can be achieved by providing expression of heterologous products or increased expression of endogenous products in plants. Alternatively, the results can be achieved by providing for a reduction of expression of one or more endogenous products, particularly enzymes or cofactors in the plant. These changes result in a change in phenotype of the transformed plant.

20           Genes of interest are reflective of the commercial markets and interests of those involved in the development of the crop. Crops and markets of interest change, and as developing nations open up world markets, new crops and technologies will emerge also. In addition, as our understanding of agronomic traits and characteristics such as yield and heterosis increase, the choice of genes for transformation will  
25 change accordingly. General categories of genes of interest include, for example, those genes involved in information, such as zinc fingers, those involved in communication, such as kinases, and those involved in housekeeping, such as heat shock proteins. More specific categories of transgenes, for example, include genes encoding important traits for agronomics, insect resistance, disease resistance,  
30 herbicide resistance, sterility, grain characteristics, and commercial products. Genes of interest include, generally, those involved in oil, starch, carbohydrate, or nutrient metabolism as well as those affecting kernel size, sucrose loading, and the like.

Agronomically important traits such as oil, starch, and protein content can be genetically altered in addition to using traditional breeding methods. Modifications include increasing content of oleic acid, saturated and unsaturated oils, increasing levels of lysine and sulfur, providing essential amino acids, and also modification of starch. Hordothionin protein modifications are described in U.S. Application Serial Nos. 08/838,763, filed April 10, 1997; U.S. Patent Nos. 5,703,049, 5,885,801, and 5,885,802; herein incorporated by reference. Another example is lysine and/or sulfur rich seed protein encoded by the soybean 2S albumin described in U.S. Patent No. 5,850,016, and the chymotrypsin inhibitor from barley, described in Williamson *et al.* (1987) *Eur. J. Biochem.* 165:99-106, the disclosures of which are herein incorporated by reference.

Derivatives of the coding sequences can be made by site-directed mutagenesis to increase the level of preselected amino acids in the encoded polypeptide. For example, the gene encoding the barley high lysine polypeptide (BHL) is derived from barley chymotrypsin inhibitor, U.S. Application Serial No. 08/740,682, filed November 1, 1996, and PCT/US97/20441, filed October 31, 1997, the disclosures of which are herein incorporated by reference. Other proteins include methionine-rich plant proteins such as from sunflower seed (Lilley *et al.* (1989) *Proceedings of the World Congress on Vegetable Protein Utilization in Human Foods and Animal Feedstuffs*, ed. Applewhite (American Oil Chemists Society, Champaign, Illinois), pp. 497-502; herein incorporated by reference); corn (Pedersen *et al.* (1986) *J. Biol. Chem.* 261:6279; Kiriara *et al.* (1988) *Gene* 71:359; both of which are herein incorporated by reference); and rice (Musumura *et al.* (1989) *Plant Mol. Biol.* 12:123, herein incorporated by reference). Other agronomically important genes encode latex, Flourey 2, growth factors, seed storage factors, and transcription factors.

Insect resistance genes may encode resistance to pests that have great yield drag such as rootworm, cutworm, European Corn Borer, and the like. Such genes include, for example, *Bacillus thuringiensis* toxic protein genes (U.S. Patent Nos. 5,366,892; 5,747,450; 5,736,514; 5,723,756; 5,593,881; and Geiser *et al.* (1986) *Gene* 48:109); lectins (Van Damme *et al.* (1994) *Plant Mol. Biol.* 24:825); and the like.

Genes encoding disease resistance traits include detoxification genes, such as against fumonisin (U.S. Patent No. 5,792,931); avirulence (avr) and disease

resistance (R) genes (Jones *et al.* (1994) *Science* 266:789; Martin *et al.* (1993) *Science* 262:1432; and Mindrinos *et al.* (1994) *Cell* 78:1089); and the like.

Herbicide resistance traits may include genes coding for resistance to herbicides that act to inhibit the action of acetolactate synthase (ALS), in particular the sulfonylurea-type herbicides (e.g., the acetolactate synthase (ALS) gene  
5 containing mutations leading to such resistance, in particular the S4 and/or Hra mutations), genes coding for resistance to herbicides that act to inhibit action of glutamine synthase, such as phosphinothricin or basta (e.g., the bar gene), or other such genes known in the art. The *bar* gene encodes resistance to the herbicide basta,  
10 the *nptII* gene encodes resistance to the antibiotics kanamycin and geneticin, and the ALS-gene mutants encode resistance to the herbicide chlorsulfuron.

Sterility genes can also be encoded in an expression cassette and provide an alternative to physical detasseling. Examples of genes used in such ways include male tissue-preferred genes and genes with male sterility phenotypes such as QM,  
15 described in U.S. Patent No. 5,583,210. Other genes include kinases and those encoding compounds toxic to either male or female gametophytic development.

The quality of grain is reflected in traits such as levels and types of oils, saturated and unsaturated, quality and quantity of essential amino acids, and levels of cellulose. In corn, modified hordothionin proteins, described in U.S. Application  
20 Serial No. 08/838,763 (filed April 10, 1997) and U.S. Patent Nos. 5,703,049, 5,885,801, and 5,885,802, provide descriptions of modifications of proteins for desired purposes.

Commercial traits can also be encoded on a gene or genes that could increase for example, starch for ethanol production, or provide expression of proteins.  
25 Another important commercial use of transformed plants is the production of polymers and bioplastics such as described in U.S. Patent No. 5,602,321. Genes such as  $\beta$ -Ketothiolase, PHBase (polyhydroxybutyrate synthase), and acetoacetyl-CoA reductase (see Schubert *et al.* (1988) *J. Bacteriol.* 170:5837-5847) facilitate expression of polyhydroxyalkanoates (PHAs). Exogenous products include plant  
30 enzymes and products as well as those from other sources including procaryotes and other eukaryotes. Such products include enzymes, cofactors, hormones, and the like. The level of proteins, particularly modified proteins having improved amino acid

distribution to improve the nutrient value of the plant, can be increased. This is achieved by the expression of such proteins having enhanced amino acid content.

The following examples are offered by way of illustration and not by way of limitation.

5

## EXPERIMENTAL

### Example 1. The Isolation of Sunflower Disease Resistant Genes and the LOX and SCIP-1 Promoter Sequences.

10

#### Materials and Methods

##### *Plant Material*

Sunflower plants were grown in the greenhouse and growth chamber. The sunflower line SMF 3 and oxox-transgenic sunflower were used for an RNA profiling study by CuraGen. Sunflower pathogen, *Sclerotinia sclerotiorum* was maintained on PDA plate at 20 °C in dark.

15

##### *Preparation of Total RNA for RNA Profiling Study and Northern Analysis*

Plant materials were ground in liquid nitrogen and total RNA was extracted by the Tri-agent Method. For each RNA profiling study, RNA samples from 6-week-old sunflower leaves and stems of transgenic sunflower plants expressing a wheat oxalate oxidase gene (oxox) were compared with RNA samples from the non-transformed parent sunflower line SMF3. Total RNA (20 ug) was separated in a 1% agarose gel containing formaldehyde. Ethidium bromide was included to verify equal loading of RNA. After transfer onto Hybond N+ membrane, the blots were hybridized with <sup>32</sup>P-labelled rhoGAP, LOX, ADH, or SCIP-1 cDNA probes. A duplicate blot was hybridized with a ribosomal 18S RNA probe as a control. Hybridization and washing conditions were performed according to Church and Gilbert.

20

25

##### *RNA Profiling*

Differences in the expression of specific genes between sunflower plants expressing a wheat oxalate oxidase gene and the sunflower line SMF3 were determined using gene expression profiling. Total RNA was analyzed using the gene

30

expression profiling process (GeneCalling®) as described in U.S. Patent No. 5,871,697, herein incorporated by reference.

*Isolation of Full-Length or Flanking Sequences by PCR Amplification of cDNA Ends*

- 5       The four cDNAs of the present invention were isolated by using RNA profiling and PCR-based technologies. RNA profiling studies were conducted through the collaboration with CuraGen Corp. The sequence information generated by the CuraGen study was used to design gene specific primers to amplifying both 3' and/or 5' end regions of the target genes using PCR-based RACE technology.
- 10       *Sclerotinia*-infected and oxox-induced cDNA libraries or cDNAs made by the Marathon cDNA Amplification Kit (Clontech) were used as template for PCR amplification. To facilitate cloning full-length cDNAs from initial cloned regions, a pair of 28 bp vector primers were designed that flanked the cDNAs on both ends (3' and 5') of the pBS vector. Amplification of either the 5' or 3' end of the cDNA was
- 15       accomplished using a vector primer (pBS-upper or pBS-lower) paired with a gene specific primer. The 5' end of a specific gene with the intact ATG start codon was cloned and sequenced. The full-length cDNA was amplified by using a new gene specific 5' primer containing sequences upstream of the ATG start codon and a 3' primer containing vector sequences.
- 20       PCR reactions were performed in a total volume of 25 µl in 10 mM Tris-HCL, pH 8.3; 1.5 mM MgCl<sub>2</sub>; 50 mM KCl; 0.1 mM dNTPs; 0.25 µM of each primer with 0.5 units of advantage cDNA polymerase mix (Clontech) or Pwo DNA polymerase (Boehringer). Genomic DNA and/or cDNA library mixtures were used as template for PCR amplification.

25

*Isolation of Disease Inducible Promoters*

- The promoter regions of the LOX and SCIP-1 genes were isolated from sunflower genomic DNA using Universal Genomic Walker Kit (clontech) according to the manufactures instructions. Restriction digested genomic DNAs were ligated
- 30       with an adapter to construct pools of genomic DNA fragments for walking by PCR.

### *Analysis of Amplified PCR Products*

Amplified PCR fragments with the expected sizes were individually sliced out of the gel for second round PCR re-amplification. The second round of PCR was carried out using the same conditions as initial PCR. Each second round PCR product that showed a single band of the expected size was cloned into the TA vector (Clontech) according to the manufacturer's instructions. Positive clones were selected for DNA sequencing using an Applied Biosystems 373A (ABI) automated sequencer at the Nucleic Acid Analysis Facility of Pioneer Hi-Bred Int'l Inc. DNA sequence analysis was carried out with the Sequencer (3.0). Multiple-sequence alignments (Clustal W) of the DNA sequence were carried out with the Curatool (CuraGen).

### *Construction of the Sclerotinia-Infected and Resistance-Enhanced (Oxox-Induced) Sunflower cDNA Libraries*

Six-week old SMF3 sunflower plants were infected with *Sclerotinia sclerotium* by petiole inoculation with *Sclerotinia*-infected carrot plugs. Six days after infection, leaf and stem tissues were collected from infected plants for total RNA isolation. Total RNA was also isolated from 6-week-old sunflower oxox-transgenic plants (line 610255) expressing a wheat oxalate oxidase gene. Our previous studies have shown that elevated levels of  $H_2O_2$ , SA, and PR1 protein were detected in oxox-transgenic plants at the 6-week stage and that the plants showed more resistance to *Sclerotinia* infection. The mRNAs were isolated using a mRNA purification kit (BRL) according to the manufacture's instructions. The cDNA libraries were constructed using the ZAP-cDNA synthesis kit and the pBluescript phagemid (Stratagene). A cDNA library mixture for PCR cloning was made of oxox-transgenic stem and *Sclerotinia*-infected leaf libraries (1:2 mix).

### *Fungal Infection and Chemical Treatments*

Sunflower plants SMF3 were planted in 4-inch pots and grown in greenhouse for first four weeks. After transfer to growth chamber, plants were maintained under 12-hour photoperiod at 22°C with an 80% relative humidity. Six-week old plants were inoculated with *Sclerotinia*-infected carrot plugs or sprayed with four different



chemicals at the given concentration. For each plant, three petioles were inoculated and wrapped with 1x2 inch parafilm. Plant tissue samples were harvested at different time points and immediately frozen in liquid nitrogen and then stored at  $-80^{\circ}\text{C}$ .

## 5 CuraGen Analysis and Database Search

An RNA profiling study identified 7 bands that were induced in oxox-transgenic and *Sclerotinia*-infected sunflower plants (Table 1).

Table 1. Summary of RNA profiling results from four sets of experiments.

10

Gene name	Fold Diff	Band Id	Stem&Leaf Oxox-48d	Leaf Oxox-48d	Leaf Infection	Stem Infection
GTPase-activating protein (rhoGAP)	+2.6	m0v0-120.8	+2.58	+3.15	0	0
Lipoxygenase	+11.8	m1n0-257.6	0	0	0	+11.82
Lipoxygenase	+6.4	w0h0-279.1	0	0	0	+6.42
Lipoxygenase	+3.4	m0r0-276.7	0	0	0	+3.37
Alcohol dehydrogenase	+7.7	i0a0-289.5	+7.69	0	0	+2.21
SCIP-1	+3.2	l0m0-273.9	0	0	+23.24	0
SCIP-1	+12.3	l0m0-94.2	0	+7.74	+12.28	0

Note: three lipoxygenase bands belong to same isolated sunflower LOX cDNA and the two SCIP-1 bands are the same cDNA.

- The CuraGen band m0v0-120.8 was induced 2 to 3 fold in oxox-transgenic plants and showed lesion mimic phenotype (Table 1). The translated product of CuraGen band m0v0-120.8 shares (amino acids 23-118 of SEQ ID NO: 2) 78% amino acid identity with an *Arabidopsis* hypothetical 23.6 kDa protein (Accession No. O81806). Band m0v0-120.8 represents amino acids 23-118 of SEQ ID NO: 2. The nucleotide sequence encoding the rhoGAP polypeptide is shown in SEQ ID NO: 1.
- The sunflower rhoGAP clone is 824 bp long with an ORF from nucleotides 35 to nucleotide 637 from the 5' end. It encodes a 201 amino acid protein (SEQ ID NO: 2) with a molecular weight of 23.4 kDa and pI of 8.1. A BLASTX (version 2.0) search indicated that the sunflower rhoGAP-1 protein has sequence homology (about 40%) with human p50rhoGAP (Accession No. Z23024) and also shares sequence homology with the *Lotus japonicus* rhoGAP (Accession No. AF064787).

Three of the CuraGen bands (m1n0-257.6, w0h0-279.1 and m0r0-276.7) were identified as lipoxygenase homologues by blast search of Genbank. The bands were

induced 3.37- to 11.82- fold higher in *Sclerotinia*-infected samples compared with uninfected control samples. The sunflower LOX nucleotide sequence is shown in SEQ ID NO: 3. The LOX ORF extends from nucleotide 881 to nucleotide 3583. The LOX ORF encodes a 901 amino acid protein (SEQ ID NO: 4) with a predicted  
 5 molecular weight of 101.37 kDa and a pI of 5.61. The sunflower LOX protein shares homology to other LOX genes from potato (Accession No. 034370), tomato (Accession No. Q96573), cowpea (Accession No. P93698), *Arabidopsis* (Accession No. LOXC\_ARATH) and rice (Accession No. LOXC\_ORYSA).

The CuraGen band i0a0-289.5 was induced 7.7-fold higher in *Sclerotinia*-  
 10 infected samples compared with uninfected control samples. PCR-based cloning of this band allowed the isolation of a cDNA encoding a 381 amino acid polypeptide, which has 85-95% amino acid identity with known plant ADH sequences from garden lettuce (Accession No. Q40249), potato (Accession No. P14674), tomato (Accession No. P28032), apple (Accession No. P48977), and maize (Accession No. P00333).

15 The CuraGen bands l0m0-273.9 and l0m0-94.2 were induced 23.2- and 12.3-fold higher in *Sclerotinia*-infected samples compared with uninfected control samples, respectfully. PCR-based cloning of this band allowed two partial cDNA clones to be isolated. The full length cDNA of SCIP-1 encodes a 168 amino acid polypeptide. A BLASTP (version 2.0) sequence alignment showed that SCIP-1 has about 41%  
 20 identity from amino acid 10-167 with a conserved protein from *Methanobacterium thermoautotrophicum* (Accession No. O26373); about 34% identity from amino acid residues 1-163 with a hypothetical 20.7 kDa protein from *Pyrococcus abyssi* (Accession No. CAB50064), about 35% identity from amino acid residues 13-159 with a conserved hypothetical protein from *Archaeoglobus fulgidus* (Accession No. O28575), about 36% identity with a hypothetical 16.5 kDa protein from *Chlamydia trachomatis* (Accession No. O84741), about 36% identity from amino acid residues  
 25 21-163, about 36% identity from amino acid 21 to 163 with a long hypothetical protein from *Pyrococcus horikoshii* (Accession No. O58984), about 35% identity with a hypothetical 18.9 kDa protein from *Aquifex aeolicus* (Accession No. O67293),  
 30 about 35% identity from amino acid residues 21-167 with a hypothetical 17.1 kDa protein from *E. coli* (Accession No. P12994), and about 35% identity from amino acid residues 8-135 with a hypothetical 19.5 kDa protein from *E. coli* (Accession No. P77368).

As discussed further in example 2, the transcript encoding SCIP-1 was up-regulated in infected tissue as compared to control tissue. The accumulation of SCIP-1 in lesion mimic and infected sunflower plants suggests that the polypeptide may be involved in the plant defense response to *Sclerotinia* and other pathogens.

5           Position specific iterative BLAST (PSI-BLAST)

(<http://www.ncbi.nlm.nih.gov/Education/BLASTinfo/psi1.html>) was performed on the SCIP-1 sequence of the invention. PSI-BLAST refers to a feature of BLAST 2.0 in which a profile (or position specific scoring matrix, PSSM) is constructed (automatically) from a multiple alignment of the highest scoring hits in an initial  
10   BLAST search. The PSSM is generated by calculating position-specific scores for each position in the alignment. Highly conserved positions receive high scores and weakly conserved positions receive scores near zero. The profile is used to perform a second (BLAST search and the results of each "iteration" used to refine the profile. The second profile is used to perform another BLAST search and so on, until the  
15   profile no longer brings back new information and is said to have "converged". This iterative searching strategy results in increased sensitivity. PSI-BLAST searches thus can identify subtle homologies to annotated entries in the database. PSI-BLAST is an important tool for predicting both biochemical activities and function from sequence relationships.

20           A search was set up using the Lion BioSCOUT implementation of the PSI-Blast algorithm. The query was the SCIP-1 polypeptide sequence, and the NR database provided with Lion BioScout was searched. Parameters were set to match as closely as possible the defaults recommended by NCBI.

25           The best PSI-Blast hits found in the database for SCIP-1 were the group of archeal and eubacterial hypothetical proteins found in earlier BLASTP searches. As the PSI-Blast search progressed, however, another group of proteins entered the alignments. Table 2 shows the 14 best hits from the PSI-Blast search other than the hypothetical proteins. It is notable that the areas of alignment are found across the entire length of these proteins (data not shown).

30           The proteins found in the PSI-Blast search fall into a class of flowering-related plant protein (CEN and others), as well as some phosphatidylethanolamine-binding proteins (PEBP). The CEN-related proteins are known to be related to a class of phosphatidylethanolamine-binding proteins (PEBP). The alignment of the SCIP-1

protein with both the CEN proteins and PEBP suggests that SCIP-1 may be related to this class of PEBP proteins. By analogy to other reported PEBP-type proteins, SCIP-1 may play a role in signaling, in membrane transduction, or in the regulation of cell death.

5

Table 2. Selected hits from PSI-Blast search with SCIP-1 against the NR protein database. From the entire set of PSI-Blast results, the 14 best hits were selected that were not from bacteria, and that had functional annotations associated with the molecule.

10

Hit (Accession Numbers)	PSI-Blast expectation	Description	Reference
trembl AB017525 AB017525_1	1e-29	gene: "BNTFL1-1"; <i>Brassica napus</i> BNTFL1-1 gene, complete cds	Mimida <i>et al.</i> (1999) <i>Plant Science</i> 142:155-162
swiss P54186 D1 ON CVO	1e-29	D1 PROTEIN (FRAGMENT) from <i>Onchocerca volvulus</i> (nematode)	Erttmann <i>et al.</i> (1996) <i>Gene</i> 174:203-7
trembl AB017528 AB017528_1	3e-29	gene: "BRTFL1-1"; <i>Brassica rapa</i> BRTFL1-1 gene, complete cds	Mimida <i>et al.</i> (1999) <i>Plant Science</i> 142:155-162
trembl AF145261 AF145261_1	1e-28	gene: "CET4"; product: "CEN-like protein 4"; <i>Nicotiana tabacum</i> CEN-like protein 4 (CET4) mRNA, complete cds.	Amaya <i>et al.</i> (1999) <i>Plant Cell</i> 11:1405-1417
trembl D87130 ATD130_1	2e-28	gene: "terminal flower1"; product: "terminal flower1"; <i>Arabidopsis thaliana</i> DNA for terminal flower1, complete cds.	Ohshima <i>et al.</i> (1997) <i>Mol Gen Genet</i> 254:186-94
trembl U84140 U84140_1	4e-28	gene: "sp"; product: "self-pruning protein"; <i>Lycopersicon esculentum</i> self-pruning protein (sp) mRNA, complete cds	Pnueli <i>et al.</i> (1989) <i>Development</i> 125:1979-1989
trembl AB027456 AB027456_1	1e-27	gene: "CiFT"; <i>Citrus unshiu</i> CiFT mRNA, complete cds.	Kobayashi <i>et al.</i> (1999) <i>Science</i> 286:1960-2
swiss O16264 PBPH CAEEL	3e-27	PHOSPHATIDYLETHANOLA MINE-BINDING PROTEIN HOMOLOG F40A3.3.	Unpublished
trembl AB027506 AB027506_1	6e-27	gene: "TSF"; product: "TSF"; <i>Arabidopsis thaliana</i> TSF (TWIN SISTER OF FT) mRNA, complete cds	Kobayashi <i>et al.</i> (1999) <i>Science</i> 286:1960-2
swiss Q41261 CEN A NTMA .	3e-26	CENTRORADIALIS PROTEIN. [ <i>Antirrhinum</i> =snapdragons, mRNA, 929 nt].	Bradley <i>et al.</i> (1996) <i>Nature</i> 379:791-7 and Bradley <i>et al.</i> (1997) <i>Science</i> 275:80-83.
trembl AB027504 AB027504_1	5e-26	gene: "FT"; product: "FT"; <i>Arabidopsis thaliana</i> FT (FLOWERING LOCUS T) mRNA, complete cds	Kobayashi <i>et al.</i> (1999) <i>Science</i> 286:1960-2

trembl AF159882 AF159882_1	8e-26	gene: "Fdr2"; product: "Cen-like protein FDR2"; <i>Oryza sativa</i> Cen-like protein FDR2 (Fdr2) mRNA, complete cds	Unpublished
trembl AB024712 AB024712_1	1e-24	gene: "ATC"; <i>Arabidopsis thaliana</i> ATC (centroradialis) gene, complete cds, strain: Landsberg	Unpublished
trembl D16111 HSHR PBP_1	1e-23	product: "human homologue of rat phosphatidylethanolamine binding protein"; Human mRNA for human homologue of rat phosphatidylethanolamine binding protein, complete cds	Hori <i>et al.</i> (1994) <i>Gene</i> 140:293-4

**Example 2: Northern Analysis of mRNA Levels of Disease Resistance Genes Following Biotic and Abiotic Stresses**

5           The expression of many plant defense genes are induced by biotic and abiotic stresses. Salicylic acid (SA), Jasmonic acid (JA), and H<sub>2</sub>O<sub>2</sub> have been implicated in playing a central role of plant disease resistance and systemic acquired resistance. Oxalic acid (OA), a compound produced by *Sclerotinia* and many other fungal pathogens *in planta*, plays an important role in the disease infection process.

10           The expression of LOX, rhoGAP, SCIP-1, and ADH mRNA in the presence of compounds known to induce either systemic acquired resistance or disease response was determined. Six-week-old sunflower leaves were sprayed until runoff with 5 mM SA, 45 μM JA, 5 mM of oxalic acid, and 5 mM H<sub>2</sub>O<sub>2</sub>. Leaf samples from chemical treated plants were collected at 0, 6, 12, and 24 hours after foliar application.

15           Northern analysis indicated that there was a significant increase in the steady-state levels of ADH mRNA in SA and H<sub>2</sub>O<sub>2</sub> treated leaves (data not shown). The highest level of ADH mRNA expression was detected at 6 hours after application of H<sub>2</sub>O<sub>2</sub> and 12 hours following the administration of SA. There was about a 2-to 3-fold increase in rhoGAP transcripts in response to both the H<sub>2</sub>O<sub>2</sub> and OA application  
20 (data not shown). The mRNA levels of SCIP-1 increased upon treatment with OA.

          JA, a product of the LOX pathway, significantly induced the steady-state levels of LOX mRNA. Therefore, LOX mRNA seems to be controlled by a positive feedback loop. In contrast, foliar application of SA and H<sub>2</sub>O<sub>2</sub> repressed the expression of LOX mRNA in sunflower. Northern and RNA profiling results  
25 revealed that sunflower LOX mRNA was elevated by *Sclerotinia* infection and oxalic acid, a pathogenic factor produced by the fungus (data not shown).

The effects of *Sclerotinia* infection and oxox on SCIP-1, rhoGAP, and ADH mRNA levels was also determined. RNA levels in *Sclerotinia*-infected sunflower and oxox-transgenic sunflower plants were determined using Northern analysis. RNA was isolated from leaves from 6-week-old non-transformed SMF3 plants and from 6-week-old oxox-transgenic plants. Steady state levels of SCIP-1 mRNA significantly increased in 6-week-old leaf tissue from the oxox-transgenic plants as compared to the control SMF3 leaf samples. RNA was also isolated from stem tissue from 6-week-old non-transformed SMF3 plants and from 6-week-old oxox-transgenic plants. An increase in steady state levels of SCIP-1 mRNA was not detected in the stem tissue (data not shown).

RNA was also isolated from 6-week-old SMF3 plants that were infected 5-days prior to sample collection with *Sclerotinia*. The steady state levels of SCIP-1 RNA following *Sclerotinia* infection increased significantly when compared to RNA levels from uninfected leaf tissue. No change in SCIP-1 RNA levels was seen in the stem tissue following *Sclerotinia* infection (data not shown).

Induction of SCIP-1, ADH, and rhoGAP expression in oxox-transgenic sunflower leaf and stem tissue during development was analyzed. RNA was isolated from leaf and stem tissue from SMF3 plants and oxox-transgenic sunflower plants at 4-week-old, 6-week-old, and 8-week-old stages. Northern blot analysis using RNA samples from SMF3 sunflower plants and samples from oxox-transgenic plants demonstrated SCIP-1 RNA levels increased in leaf tissues in the oxox-transgenic sunflowers. The increase in SCIP-1 RNA levels was most significant at the 8-week-old stage. No detectable induction of SCIP-1 was seen in the stem tissue.

The steady-state-level of ADH mRNA is much higher in stem than in leaf tissue. Adh expression was induced in 8-week-old oxox-transgenic leaf tissue. However, its expression was repressed in oxox-stem tissue. CuraGen QEA assay results indicate that Adh expression was induced by *Sclerotinia* infection in stem and leaf tissues (data not shown).

In the leaf tissue, rhoGAP mRNA level was induced by oxox expression at the 4-week-old stage, and then was repressed with development. In stem tissue, rhoGAP expression was slightly induced by oxox (data not shown).

LOX expression in response to wounding and the ABA signal was determined. Six-week-old sunflower plants (SMF3) were sprayed with 100  $\mu$ M ABA until the

chemical solution started to run off the leaves. For the wounding experiment, each wounded leaf was crushed by a hemostat 20 times and three leaves were treated from each plant. At each time point, six leaves were collected from two treated plants and immediately frozen in liquid N<sub>2</sub>. Total RNA was isolated and Northern analysis was performed. Northern blot analysis indicated that wounding significantly induced the steady state levels of LOX mRNA. The peak of the LOX mRNA accumulation was detected at 6 hours after wounding and high levels of LOX expression was maintained through 72 hours after initial treatment. ABA treatment showed only a slight induction of LOX expression at 12 hours after treatment (data not shown).

### Example 3: Transformation and Regeneration of Maize Transgenic Plants

The nucleotide sequences of the present invention can be used to transform sunflower, maize, or other plants using *Agrobacterium* or particle-gun methods. Examples 3, 4, and 5 provide methods for sunflower, maize, and soybean transformations.

Immature maize embryos from greenhouse donor plants are bombarded with a plasmid containing a rhoGAP, LOX, ADH or SCIP-1 nucleotide sequence operably linked to a ubiquitin promoter (Figure 1). The plasmids also contain the selectable marker gene PAT (Wohlleben *et al.* (1988) *Gene* 70:25-37) that confers resistance to the herbicide Bialaphos. Transformation is performed as follows. All media recipes are given in the Appendix. Alternatively, the plant can be transformed with a plasmid comprising the LOX or SCIP-1 promoter sequences of the invention operably linked to the nucleotide sequence encoding the GUS reporter protein (Figure 2).

### 25 Preparation of Target Tissue

The ears are surface sterilized in 30% Chlorox bleach plus 0.5% Micro detergent for 20 minutes, and rinsed two times with sterile water. The immature embryos are excised and placed embryo axis side down (scutellum side up), 25 embryos per plate, on 560Y medium for 4 hours and then aligned within the 2.5-cm target zone in preparation for bombardment.

### *Preparation of DNA*

A plasmid vector shown in Figures 1 or 2 is precipitated onto 1.1  $\mu\text{m}$  (average diameter) tungsten pellets using a  $\text{CaCl}_2$  precipitation procedure as follows:

- 5        100  $\mu\text{l}$  prepared tungsten particles in water
- 10  $\mu\text{l}$  (1  $\mu\text{g}$ ) DNA in TrisEDTA buffer (1  $\mu\text{g}$  total)
- 100  $\mu\text{l}$  2.5 M  $\text{CaCl}_2$
- 10  $\mu\text{l}$  0.1 M spermidine

- Each reagent is added sequentially to the tungsten particle suspension, while
- 10       maintained on the multitube vortexer. The final mixture is sonicated briefly and allowed to incubate under constant vortexing for 10 minutes. After the precipitation period, the tubes are centrifuged briefly, liquid removed, washed with 500 ml 100% ethanol, and centrifuged for 30 seconds. Again the liquid is removed, and 105  $\mu\text{l}$  100% ethanol is added to the final tungsten particle pellet. For particle gun
  - 15       bombardment, the tungsten/DNA particles are briefly sonicated and 10  $\mu\text{l}$  spotted onto the center of each macrocarrier and allowed to dry about 2 minutes before bombardment.

### *Particle Gun Treatment*

- 20       The sample plates are bombarded at level #4 in particle gun #HE34-1 or #HE34-2. All samples receive a single shot at 650 PSI, with a total of ten aliquots taken from each tube of prepared particles/DNA.

### *Subsequent Treatment*

- 25       Following bombardment, the embryos are kept on 560Y medium for 2 days, then transferred to 560R selection medium containing 3 mg/liter Bialaphos, and subcultured every 2 weeks. After approximately 10 weeks of selection, selection-resistant callus clones are transferred to 288J medium to initiate plant regeneration. Following somatic embryo maturation (2-4 weeks), well-developed somatic embryos
- 30       are transferred to medium for germination and transferred to the lighted culture room. Approximately 7-10 days later, developing plantlets are transferred to 272V hormone-free medium in tubes for 7-10 days until plantlets are well established. Plants are then



transferred to inserts in flats (equivalent to 2.5" pot) containing potting soil and grown for 1 week in a growth chamber, subsequently grown an additional 1-2 weeks in the greenhouse, then transferred to classic 600 pots (1.6 gallon) and grown to maturity.

Plants transformed with the plasmid shown in Figure 1 comprising the  
5 rhoGAP, LOX, ADH, or SCIP-1 nucleotide sequences are monitored and scored for altered defense response, or altered rhoGAP, LOX, ADH, or SCIP-1 activity.

Plants transformed with the plasmid shown in Figure 2 comprising the LOX or SCIP-1 promoter sequences operably linked to the GUS reporter sequences are monitored for LOX or SCIP-1 promoter activity. Following exposure to various  
10 stimuli, such as for example, *Sclerotinia* and oxalic acid, LOX promoter activity is measured using the reporter gene GUS. GUS activity in various tissues is measured by a fluorogenic assay. The fluorogenic assay determines the specific activity of  $\beta$ -glucuronidase (GUS) in various maize tissue extracts. The specific activity of the enzyme is expressed as moles of 4-methyl umbelliferone (MU) released/ $\mu$ g  
15 protein/hour. MU is produced when the enzyme (GUS) in plant cell extracts cleaves the glucuronide moiety from the 4-methyl umbelliferyl- $\beta$ -D-glucuronide (MUG) substrate.

#### *Bombardment and Culture Media*

20 Bombardment medium (560Y) comprises 4.0 g/l N6 basal salts (SIGMA C-1416), 1.0 ml/l Eriksson's Vitamin Mix (1000X SIGMA-1511), 0.5 mg/l thiamine HCl, 120.0 g/l sucrose, 1.0 mg/l 2,4-D, and 2.88 g/l L-proline (brought to volume with D-I H<sub>2</sub>O following adjustment to pH 5.8 with KOH); 2.0 g/l Gelrite (added after bringing to volume with D-I H<sub>2</sub>O); and 8.5 mg/l silver nitrate (added after sterilizing  
25 the medium and cooling to room temperature). Selection medium (560R) comprises 4.0 g/l N6 basal salts (SIGMA C-1416), 1.0 ml/l Eriksson's Vitamin Mix (1000X SIGMA-1511), 0.5 mg/l thiamine HCl, 30.0 g/l sucrose, and 2.0 mg/l 2,4-D (brought to volume with D-I H<sub>2</sub>O following adjustment to pH 5.8 with KOH); 3.0 g/l Gelrite (added after bringing to volume with D-I H<sub>2</sub>O); and 0.85 mg/l silver nitrate and 3.0  
30 mg/l bialaphos(both added after sterilizing the medium and cooling to room temperature).

Plant regeneration medium (288J) comprises 4.3 g/l MS salts (GIBCO 11117-074), 5.0 ml/l MS vitamins stock solution (0.100 g nicotinic acid, 0.02 g/l thiamine

HCL, 0.10 g/l pyridoxine HCL, and 0.40 g/l glycine brought to volume with polished D-I H<sub>2</sub>O) (Murashige and Skoog (1962) *Physiol. Plant.* 15:473), 100 mg/l myo-inositol, 0.5 mg/l zeatin, 60 g/l sucrose, and 1.0 ml/l of 0.1 mM abscisic acid (brought to volume with polished D-I H<sub>2</sub>O after adjusting to pH 5.6); 3.0 g/l Gelrite (added after bringing to volume with D-I H<sub>2</sub>O); and 1.0 mg/l indoleacetic acid and 3.0 mg/l bialaphos (added after sterilizing the medium and cooling to 60°C). Hormone-free medium (272V) comprises 4.3 g/l MS salts (GIBCO 11117-074), 5.0 ml/l MS vitamins stock solution (0.100 g/l nicotinic acid, 0.02 g/l thiamine HCL, 0.10 g/l pyridoxine HCL, and 0.40 g/l glycine brought to volume with polished D-I H<sub>2</sub>O), 0.1 g/l myo-inositol, and 40.0 g/l sucrose (brought to volume with polished D-I H<sub>2</sub>O after adjusting pH to 5.6); and 6 g/l bacto-agar (added after bringing to volume with polished D-I H<sub>2</sub>O), sterilized and cooled to 60° C.

#### Example 4: *Agrobacterium*-mediated Transformation

For *Agrobacterium*-mediated transformation of maize with rhoGAP, LOX, ADH, or SCIP-1 nucleotide sequences of the invention or a nucleotide sequence operably linked to the LOX or SCIP-1 promoter sequence of the invention, preferably the method of Zhao is employed (U.S. Patent No. 5,981,840, and PCT patent publication WO98/32326; the contents of which are hereby incorporated by reference). Briefly, immature embryos are isolated from maize and the embryos contacted with a suspension of *Agrobacterium*, where the bacteria are capable of transferring the DNA constructs of interest to at least one cell of at least one of the immature embryos (step 1: the infection step). In this step the immature embryos are preferably immersed in an *Agrobacterium* suspension for the initiation of inoculation. The embryos are co-cultured for a time with the *Agrobacterium* (step 2: the co-cultivation step). Preferably the immature embryos are cultured on solid medium following the infection step. Following this co-cultivation period an optional "resting" step is contemplated. In this resting step, the embryos are incubated in the presence of at least one antibiotic known to inhibit the growth of *Agrobacterium* without the addition of a selective agent for plant transformants (step 3: resting step). Preferably the immature embryos are cultured on solid medium with antibiotic, but without a selecting agent, for elimination of *Agrobacterium* and for a resting phase for the infected cells. Next, inoculated embryos are cultured on medium containing a

selective agent and growing transformed callus is recovered (step 4: the selection step). Preferably, the immature embryos are cultured on solid medium with a selective agent resulting in the selective growth of transformed cells. The callus is then regenerated into plants (step 5: the regeneration step), and preferably calli grown on selective medium are cultured on solid medium to regenerate the plants.

#### Example 5: Transformation and Regeneration of Sunflower Plants

The intact meristem method is used for transformation of sunflower plants and expression of the LOX, ADH, rhoGAP, or SCIP-1 nucleotide sequences as follows.

Alternatively, the same method could be used to express a nucleotide sequence of interest under the control of the LOX or SCIP-1 promoter sequence of the invention.

#### *Explant Preparation*

Seeds are dehulled and surface-sterilized for 20 minutes in a 20% Chlorox™ bleach solution with the addition of two to three drops of Tween 20 per 100 ml of solution, and then rinsed three times with distilled water. Sterilized seeds are imbibed in the dark at 26°C for 20 hours on filter paper moistened with water. Meristem explants are created by removing cotyledons and root radicle from imbibed seeds, and then culturing overnight at 26°C in the dark on 374E medium (1X MS salts, Shepards vitamins, 40 mg/l adenine sulfate, 30 g/l sucrose, 0.5 mg/l BAP, 0.25 mg/l IAA, 0.1 mg/l IAA, pH 5.6, 8g/l phytagar). Primary leaves are then removed and explants are transferred to 374M medium (374E except 12 g/l phytagar), arranged in a manner suitable for particle gun bombardment, and cultured overnight at 26°C in the dark.

#### *Preparation of DNA*

A plasmid vector comprising the rhoGAP, LOX, ADH, or SCIP-1 nucleotide sequences operably linked to a ubiquitin promoter is constructed (Figure 1). Alternatively, a plasmid vector comprising the LOX promoter or the SCIP-1 nucleotide sequence operably linked to the nucleotide sequence encoding the GUS reporter protein is constructed (Figure 1). Both of these plasmids contain a kanamycin selectable marker gene. The transformation is performed as follows.

### Transformation

Approximately 18.8 mg of 1.8  $\mu$ m tungsten particles are suspended in 150  $\mu$ l absolute ethanol, and sonicated for 2-4 seconds. After sonication, 10  $\mu$ l of the suspension is dropped on the center of the surface of a macrocarrier. Each plate of meristem explants is bombarded twice with 650 psi rupture discs in the top shelf at 26 mm of Hg helium gun vacuum, using a BioRad helium gun.

The plasmid vector shown in Figure 1 or in Figure 2 is introduced into *Agrobacterium* strain EHA 105 (see above) via freeze-thawing as described by Holsters *et al.* (1978) *Mol. Gen. Genet.* 163:181-187. Actively growing, transformed *Agrobacteria* were maintained in shaking liquid cultures using 60A medium with kanamycin (YEP, 50 mg/l kanamycin: 10 g/l yeast extract, 10 g/l bactopectone, 5 g/l NaCl, pH 7.0, 50 mg/l kanamycin). On the day before the *Agrobacterium* strain is to be used, new liquid cultures are initiated in 60A with kanamycin from the active maintenance culture. They are cultured with shaking at 26°C until they reach an optical density (OD vis = 600 nm) of about 1.0. When the cultures have established this density, they are centrifuged (6000 rpm, 5 min), the supernatant is discarded, and the pellet of bacteria is resuspended in inoculation medium (12.5mM 2-(N-morpholino) ethanesulfonic acid, 1 g/l NH<sub>4</sub>Cl, and 0.3 g/l MgSO<sub>4</sub>, at pH 5.7), to a final calculated concentration of *Agrobacteria* of 4.0 at OD 600. The particle bombarded explants are inoculated with *Agrobacterium* by first spreading the explants apart on the 374M medium, then placing a droplet of the above suspension directly onto the top of each meristem. The explants are co-cultivated on the medium for 4 days, after which the explants are transferred to 374 C medium (GBA with 1% sucrose and with no BAP, IAA, or GA<sub>3</sub>, and supplemented with 250  $\mu$ g/ml cefotaxime). The explants are cultured on this medium for about 2 weeks under 16 hours of daylight, at 26°C.

### Recovering Nodes and Plants

Following the 4 days of co-cultivation time on 374M medium, the explants are transferred to 374D (374C medium with 50 mg/l kanamycin) selection medium containing kanamycin. After 2 weeks of selection, explants with associated shoots are transferred to 374C medium and selection resistant shoots are screened using NPTII

ELISA. Positive shoots are removed for recovery by *in vitro* grafting and transformation verified by further NPTII ELISA analysis. Negative shoots are discarded. Explants with smaller shoots which could not be assayed following the 2 weeks on 374D are transferred to 374G (374E with 250 mg/l cefotaxime) for 3-4 days then back to 374C for 2 additional weeks. Assays are then done to identify positive shoots which are too small to sample in the first round and recovery initiated.

Recovered positive shoots are grafted to Pioneer sunflower hybrid *in vitro*-grown sunflower seedling rootstock. The seeds are dehulled and surface-sterilized for 20 minutes in a 20% Chlorox<sup>TM</sup> bleach solution with two to three drops of Tween20 per 100 ml total volume, and rinsed three times with distilled water. The sterilized seeds are germinated for three days on filter paper moistened with water, then transferred into "48 Medium" (one-half strength MS salts, 0.5% sucrose, 0.3% gelrite, at pH 5.0) and grown at 26°C at 26 in the dark for 3 days, then incubated at 16 hour day culture conditions. The upper portions of selected seedlings are removed, a vertical slice is made in each hypocotyl, and a transformed shoot is inserted into the vertical slice. The cut area is wrapped with parafilm, and after one week culture on the medium, the grafted plants are transferred to soil. In the first two weeks they are maintained under high humidity conditions to acclimatize to the greenhouse environment.

Transformed sectors of TO plants are identified by additional NPTII assays of the greenhouse established positive grafted shoots. After assay, non-transformed sectors are trimmed off to promote auxillary bud development and auxiliary buds from transgenic sectors are recovered so as to establish the best probability to encompass the sector of transformation in germ line cells so that the transformation event is recovered in the next generation. Seed from TO plants are collected, dehulled, surface sterilized, and germinated on filter paper wetted with water. T1 seedlings are then sampled for NPTII ELISA by removing green cotyledon pieces followed by transfer to seedling growth medium 48P (0.1X MS salts, 0.5% sucrose, pH 5.6, 0.3% gelrite). NPTII positive, actively growing T1 seedlings are transferred at the two leaf stage to soil for growth in the greenhouse. Seed from the confirmed T1 transgenics is used to grow T2 plants.

T2 seeds are planted in a greenhouse. Positive plants are screened by NPTII assay. Various plant tissues are harvested at 80-day-old stage after planting. The harvested material is put in mini-tubes, frozen and stored at  $-80^{\circ}\text{C}$ .

Plants transformed with the plasmid shown in Figure 1 comprising the  
5 rhoGAP, LOX, ADH, or SCIP-1 nucleotide sequences are monitored and scored for an altered defense response, or a modulation in rhoGAP, LOX, ADH or SCIP-1 activity.

Plants transformed with the plasmid shown in Figure 2 comprising the LOX promoter sequences operably linked to the GUS reporter sequences are monitored for  
10 LOX promoter activity. Following exposure to various stimuli that induce the LOX promoter, LOX promoter activity is measured by assaying for GUS activity. GUS activity in various tissues is measured by a fluorogenic assay. The fluorogenic assay determines the specific activity of  $\beta$ -glucuronidase (GUS) in various sunflower tissue extracts. The specific activity of the enzyme is expressed as moles of 4-methyl  
15 umbelliferone (MU) released/ $\mu\text{g}$  protein/hour. MU is produced when the enzyme (GUS) in plant cell extracts cleaves the glucuronide moiety from the 4-methyl umbelliferyl- $\beta$ -D-glucuronide (MUG) substrate.

Harvested T2 tissue samples stored at  $-80^{\circ}\text{C}$  are homogenized in 400  $\mu\text{l}$  lysis buffer (40 mM Phosphate, pH 7.0, 10 mM EDTA, 10 mM  $\beta$ -mercaptoethanol), and  
20 then centrifuged in the Jouan GR422 centrifuge for 10 minutes at 4000 rpm. The total protein concentration of the supernatant is measured using the Bio-Rad Bradford Method (Bio-RAD) with BSA as the standard protein according to manufacture's protocol. Ten  $\mu\text{l}$  of diluted supernatant (about 4  $\mu\text{g}$  of total protein) is used for the GUS activity assay. GUS activity is assayed according to Jefferson *et al.* (1987)  
25 *EMBO J.* 6: 3901-3907 using MUG as substrate.

As an alternative to the intact meristem method, the split embryonic axis method may be used as described in Malone-Schoneberg *et al.* (1994) *Plant Science* 103:193-207, in transforming sunflower plants with either the plasmid shown in Figure 3 or Figure 4 and generating T2 plants. T2 seeds are planted in a greenhouse  
30 and positive plants are screened by NPTII assay. Plant tissues are harvested at 80-day-old stage after planting. The harvested material is put in mini-tubes, frozen, and stored at  $-80^{\circ}\text{C}$ .

Example 6: Soybean Embryo Transformation

Soybean embryos are bombarded with a plasmid containing the SCIP-1, rhoGAP, LOX, or ADH sequences operably linked to a ubiquitin promoter (Figure 1) as follows. Alternatively, the soybean embryos can be bombarded with a DNA construct containing the SCIP-1 or LOX promoter operably linked to a nucleotide sequence of interest (Figure 2). To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface-sterilized, immature seeds of the soybean cultivar A2872, are cultured in the light or dark at 26°C on an appropriate agar medium for six to ten weeks. Somatic embryos producing secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos that multiplied as early, globular-staged embryos, the suspensions are maintained as described below.

Soybean embryogenic suspension cultures can be maintained in 35 ml liquid media on a rotary shaker, 150 rpm, at 26°C with fluorescent lights on a 16:8 hour day/night schedule. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 ml of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein *et al.* (1987) *Nature* (London) 327:70-73, U.S. Patent No. 4,945,050). A Du Pont Biolistic PDS1000/HE instrument (helium retrofit) can be used for these transformations.

A selectable marker gene that can be used to facilitate soybean transformation is a transgene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell *et al.* (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz *et al.* (1983) *Gene* 25:179-188), and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The expression cassette comprising the DNA construct can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

To 50 µl of a 60 mg/ml 1 µm gold particle suspension is added (in order): 5 µl DNA (1 µg/µl), 20 µl spermidine (0.1 M), and 50 µl CaCl<sub>2</sub> (2.5 M). The particle preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 µl

70% ethanol and resuspended in 40  $\mu$ l of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five microliters of the DNA-coated gold particles are then loaded on each macro carrier disk.

Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi, and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post-bombardment with fresh media containing 50 mg/ml hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post-bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or regenerated into whole plants by maturation and germination of individual somatic embryos.


All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.



INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL


<b>A.</b> The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 10, lines 23, 27 and 31	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 30 June 1999 (30.06.99)	Accession Number PTA-284
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 11, lines 11, 13, 16, 18 and 19; Page 78, lines 9 and 13; Page 79, lines 4 and 6; Page 80, lines 8 and 10; Page 81, lines 11 and 13; Page 83, lines 11 and 13	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indicators are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL


<b>A.</b> The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 10, lines 23, 28 and 31	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 30 June 1999 (30.06.99)	Accession Number PTA-285
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 11, lines 11, 13, 17-19; Page 78, lines 9 and 13; Page 79, lines 4 and 6; Page 80, lines 8 and 10; Page 81, lines 11 and 13; Page 83, lines 11 and 13	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indicators are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL

<b>A.</b> The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 10, lines 24 and 31	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 30 June 1999 (30.06.99)	Accession Number PTA-286
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 11, lines 11 and 13; Page 78, line 9; Page 79, line 4; Page 80, line 8; Page 81, line 11; Page 83, line 11	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indicators are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL

<b>A.</b> The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 10, lines 25 and 31	
<b>B. IDENTIFICATION OF DEPOSIT</b>	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 30 June 1999 (30.06.99)	Accession Number PTA-287
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable)	
This information is continued on an additional sheet <input type="checkbox"/>	
Page 11, lines 12 and 13; Page 78, line 9; Page 79, line 4; Page 80, line 8; Page 81, line 11; Page 83, line 11	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indicators are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	
<input checked="" type="checkbox"/> This sheet was received with the international application	
Authorized officer Arlene Brindle	

For International Bureau use only	
<input type="checkbox"/> This sheet was received with the International Bureau on:	
Authorized officer	

INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL

<b>A.</b> The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 10, lines 25 and 31	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 30 June 1999 (30.06.99)	Accession Number PTA-288
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 11, lines 12 and 13; Page 78, line 10; Page 79, line 4; Page 80, line 8; Page 81, line 11; Page 83, line 11	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indicators are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer Arlene Bringle

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL


A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 11, lines 4, 12 and 14	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 20 August 1999 (20.08.99)	Accession Number PTA-559
C. ADDITIONAL INDICATIONS (leave blank if not applicable)	
This information is continued on an additional sheet <input type="checkbox"/>	
Page 82, line 5; Page 84, lines 4 and 17; Page 85, lines 1 and 15	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indicators are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer Arlene Bringle

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL

<b>A.</b> The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 11, lines 4, 12 and 15	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution <b>American Type Culture Collection</b>	
Address of depository institution (including postal code and country) <b>10801 University Blvd. Manassas, VA 20110-2209 US</b>	
Date of deposit <b>26 April 2000 (26.04.00)</b>	Accession Number <b>PTA-1721</b>
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 82, line 6; Page 84, lines 5 and 18; Page 85, lines 2 and 16	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indicators are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the International application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

## THAT WHICH IS CLAIMED:

1. An isolated polypeptide selected from the group consisting of:
  - (a) a polypeptide comprising an amino acid sequence set forth in  
5 SEQ ID NO: 2, 4, 7, or 9;
  - (b) a polypeptide encoded by a nucleotide sequence comprising the  
sequence set forth in SEQ ID NOS: 1, 3, 6, or 8;
  - (c) a polypeptide comprising an amino acid sequence encoded by a  
nucleotide sequence deposited as Accession No. PTA-284, PTA-285, PTA-286, PTA-  
10 287, or PTA-288;
  - (d) a polypeptide comprising an amino acid sequence encoded by a  
nucleotide sequence obtained from the overlapping clones deposited as Accession No.  
PTA-284 and PTA-285;
  - (e) a polypeptide encoded by a nucleotide sequence that hybridizes  
15 under stringent conditions to a nucleotide sequence comprising the sequences set forth  
in SEQ ID NOS: 1, 3, 6, or 8;
  - (f) an amino acid sequence having at least 60% sequence identity  
to the sequence set forth in SEQ ID NO: 4, wherein said sequence has LOX-like  
activity;
  - 20 (g) an amino acid sequence having at least 60% sequence identity  
to the sequence set forth in SEQ ID NO: 9, wherein said sequence has SCIP-1-like  
activity;
  - (h) an amino acid sequence having at least 70% sequence identity  
to the sequence set forth in SEQ ID NO: 2, wherein said sequence has rhoGAP-like  
25 activity; and,
  - (i) an amino acid sequence having at least 90% sequence identity  
to the sequence set forth in SEQ ID NO: 7, wherein said sequence has ADH-like  
activity.
- 30 2. An isolated nucleic acid molecule selected from the group consisting  
of:
  - (a) a nucleic acid molecule comprising a nucleotide sequence set  
forth in SEQ ID NO: 1, 3, 6 or 8;



- (b) a nucleic acid molecule comprising a nucleotide sequence encoding an amino acid sequence set forth in SEQ ID NO: 2, 4, 7 or 9;
- (c) a nucleic acid molecule comprising a nucleotide sequence deposited as Accession Nos. PTA-284, PTA-285, PTA-286, PTA-287, or PTA-288;
- 5 (d) a nucleic acid molecule comprising a nucleotide sequence obtained from the overlapping clones deposited as Accession No. PTA-284 and PTA-285;
- (e) a nucleic acid molecule comprising an antisense sequence corresponding to a sequence of a), b), c), or d);
- 10 (f) a nucleic acid molecule comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequences of a), b), c), d), or e), wherein said sequence encodes a polypeptide having rhoGAP-, LOX-, SCIP-1, or ADH-like activity;
- (g) a nucleic acid molecule comprising a nucleotide sequence
- 15 having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 1, wherein said sequence encodes a polypeptide having rhoGAP-like activity;
- (h) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 3, wherein said sequence encodes a polypeptide having LOX-like activity;
- 20 (i) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 6, wherein said sequence encodes a polypeptide having ADH-like activity; and,
- (j) a nucleic acid molecule comprising a nucleotide sequence
- 25 having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 8, wherein said sequence encodes a polypeptide having SCIP-1-like activity.

3. A DNA construct comprising a nucleotide sequence of claim 2, wherein said nucleotide sequence is operably linked to a promoter that drives expression in a host cell.

30

4. A cell having stably incorporated into its genome at least one DNA construct comprising a nucleotide sequence operably linked to a promoter that drives

expression in said cell, wherein said nucleotide sequence is selected from the group consisting of:

- (a) a nucleic acid molecule comprising a nucleotide sequence set forth in SEQ ID NO: 1, 3, 6 or 8;
- 5 (b) a nucleic acid molecule comprising a nucleotide sequence encoding an amino acid sequence set forth in SEQ ID NO: 2, 4, 7 or 9;
- (c) a nucleic acid molecule comprising a nucleotide sequence deposited as Accession Nos. PTA-284, PTA-285, PTA-286, PTA-287, or PTA-288;
- (d) a nucleic acid molecule comprising a nucleotide sequence  
10 obtained from the overlapping clones deposited as Accession No. PTA-284 and PTA-285;
- (e) a nucleic acid molecule comprising an antisense sequence corresponding to a sequence of a), b), c), or d);
- (f) a nucleic acid molecule comprising a nucleotide sequence that  
15 hybridizes under stringent conditions to the nucleotide sequences of a), b), c), d), or e), wherein said sequence encodes a polypeptide having rhoGAP-, LOX-, SCIP-1, or ADH-like activity;
- (g) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 1,  
20 wherein said sequence encodes a polypeptide having rhoGAP-like activity;
- (h) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 3, wherein said sequence encodes a polypeptide having LOX-like activity;
- (i) a nucleic acid molecule comprising a nucleotide sequence  
25 having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 6, wherein said sequence encodes a polypeptide having ADH-like activity; and,
- (j) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 8, wherein said sequence encodes a polypeptide having SCIP-1-like activity.

30

5. The cell of claim 4, wherein said cell is a plant cell.

6. A transformed plant having stably incorporated into its genome at least one DNA construct comprising a nucleotide sequence operably linked to a heterologous promoter that drives expression in a plant cell, wherein said nucleotide sequence is selected from the group consisting of:

- 5 (a) a nucleic acid molecule sequence comprising a nucleotide sequence set forth in SEQ ID NO: 1, 3, 6 or 8;
- (b) a nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 2, 4, 7 or 9;
- 10 (c) a nucleic acid molecule comprising a nucleotide sequence deposited as Accession Nos. PTA-284, PTA-285, PTA-286, PTA-287, or PTA-288;
- (d) a nucleic acid molecule comprising a nucleotide sequence obtained from the overlapping clones deposited as Accession No. PTA-284 and PTA-285;
- 15 (e) a nucleic acid molecule comprising an antisense sequence corresponding to a sequence of a), b), c), or d);
- (f) a nucleic acid molecule comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequences of a), b), c), d), or e), wherein said sequence encodes a polypeptide having rhoGAP-, LOX-, SCIP-1, or
- 20 ADH-like activity;
- (g) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 1, wherein said sequence encodes a polypeptide having rhoGAP-like activity;
- (h) a nucleic acid molecule comprising a nucleotide sequence
- 25 having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 3, wherein said sequence encodes a polypeptide having LOX-like activity;
- (i) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 6, wherein said sequence encodes a polypeptide having ADH-like activity; and,
- 30 (j) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 8, wherein said sequence encodes a polypeptide having SCIP-1-like activity.

7. The DNA construct of claim 5 wherein said promoter is selected from the group consisting of:

(a) a nucleic acid molecule comprising a nucleotide sequence having the sequence set forth in SEQ ID NO: 5 or SEQ ID NO: 10;

5 (b) a nucleic acid molecule deposited as Accession No. PTA-559 or PTA-1721;

(c) a nucleic acid molecule comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequences of a) or b) wherein said sequence retains the ability to regulate transcription; and,

10 (d) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to a sequence set forth in SEQ ID NOS: 5 or 10, wherein said sequence retains the ability to regulate transcription.

15 8. The plant of claim 6, wherein said promoter is a constitutive promoter.

9. The plant of claim 6, wherein said promoter is a tissue-preferred promoter.

20 10. The plant of claim 6, wherein said promoter is an inducible promoter.

11. The plant of claim 10, wherein said promoter is a pathogen-inducible promoter.

25 12. The plant of claim 6, wherein said plant is a monocot.

13. The plant of claim 12, wherein said monocot is maize, wheat, rice, barley, sorghum, or rye.

30 14. The plant of claim 6, wherein said plant is a dicot.

15. The transformed seed of the plant of claim 6.

16. A method for enhancing a plant defense response, said method comprising stably introducing into the genome of a plant at least one DNA construct comprising a nucleotide sequence operably linked to a heterologous promoter that drives expression in a plant cell, wherein said nucleotide sequence is selected from the group consisting of:
- 5 (a) a nucleic acid molecule sequence comprising a nucleotide sequence set forth in SEQ ID NO: 1, 3, 6 or 8;
- (b) a nucleic acid molecule comprising a nucleotide sequence encoding an amino acid sequence set forth in SEQ ID NO: 2, 4, 7 or 9;
- 10 (c) a nucleic acid molecule comprising a nucleotide sequence deposited as Accession Nos. PTA-284, PTA-285, PTA-286, PTA-287, or PTA-288;
- (d) a nucleic acid molecule comprising a nucleotide sequence obtained from the overlapping clones deposited as Accession No. PTA-284 and PTA-285;
- 15 (e) a nucleic acid molecule comprising an antisense sequence corresponding to a sequence of a), b), c), or d);
- (f) a nucleic acid molecule comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequences of a), b), c), d), or e), wherein said sequence encodes a polypeptide having rhoGAP-, LOX-, SCIP-1, or
- 20 ADH-like activity;
- (g) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 1, wherein said sequence encodes a polypeptide having rhoGAP-like activity;
- (h) a nucleic acid molecule comprising a nucleotide sequence
- 25 having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 3, wherein said sequence encodes a polypeptide having LOX-like activity;
- (i) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 6, wherein said sequence encodes a polypeptide having ADH-like activity; and,
- 30 (j) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to the sequence set forth in SEQ ID NO: 8, wherein said sequence encodes a polypeptide having SCIP-1-like activity.

17. An isolated nucleotide sequence selected from the group consisting of:
- (a) a nucleic acid molecule comprising a nucleotide sequence having the sequence set forth in SEQ ID NO: 5 or SEQ ID NO: 10;
  - (b) a nucleic acid molecule deposited as Accession No. PTA-559 or PTA-1721;
  - (c) a nucleic acid molecule comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequences of a) or b) wherein said sequence retains the ability to regulate transcription; and,
  - (d) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to a sequence set forth in SEQ ID NOS: 5 or 10, wherein said sequence retains the ability to regulate transcription.
18. A DNA construct comprising a promoter operably linked to a nucleotide sequence, wherein said promoter is selected from the group consisting of:
- (a) a nucleic acid molecule comprising a nucleotide sequence having the sequence set forth in SEQ ID NO: 5 or SEQ ID NO: 10;
  - (b) a nucleic acid molecule deposited as Accession No. PTA-559 or PTA-1721;
  - (c) a nucleic acid molecule comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequences of a) or b) wherein said sequence retains the ability to regulate transcription; and,
  - (d) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to a sequence set forth in SEQ ID NOS: 5 or 10, wherein said sequence retains the ability to regulate transcription.
19. An expression vector comprising the DNA construct of claim 18.
20. A plant having stably incorporated into its genome at least one DNA construct comprising a nucleotide sequence encoding a protein of interest operably linked to a promoter, wherein said promoter is selected from the group consisting of:
- (a) a nucleic acid molecule comprising a nucleotide sequence having the sequence set forth in SEQ ID NO: 5 or SEQ ID NO: 10;

(b) a nucleic acid molecule deposited as Accession No. PTA-559 or PTA-1721;

(c) a nucleic acid molecule comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequences of a) or b) wherein  
5 said sequence retains the ability to regulate transcription; and,

(d) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to a sequence set forth in SEQ ID NOS: 5 or 10, wherein said sequence retains the ability to regulate transcription.

10 21. A cell having stably incorporated into its genome at least one DNA construct comprising a nucleotide sequence encoding a protein of interest operably linked to a promoter, wherein said promoter is selected from the group consisting of:

(a) a nucleic acid molecule comprising a nucleotide sequence having the sequence set forth in SEQ ID NO: 5 or SEQ ID NO: 10;

15 (b) a nucleic acid molecule deposited as Accession No. PTA-559 or PTA-1721;

(c) a nucleic acid molecule comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequences of a) or b) wherein said sequence retains the ability to regulate transcription; and,


20 (d) a nucleic acid molecule comprising a nucleotide sequence having at least 60% sequence identity to a sequence set forth in SEQ ID NOS: 5 and 10, wherein said sequence retains the ability to regulate transcription.

25 22. The cell of claim 21, wherein said cell is a plant cell.

23. A method of regulating the expression of a nucleotide sequence of interest, said method comprising stably incorporating in the genome of a plant cell a nucleotide sequence of interest operably linked to a promoter comprising the nucleotide sequence of claim 17.


30

24. The method of claim 23, further comprising contacting said cell with a stimuli that induces expression of said nucleotide sequence of interest.



**FIG. 1.**





**FIG. 2.**

## SEQUENCE LISTING

<110> Bidney, Dennis  
 Duvick, Jon  
 Hendrick, Carol  
 Hu, Xu  
 Lu, Guihua  
 Crasta, Oswald

<120> Sunflower RhoGAP, LOX, ADH and SCIP -  
 Polynucleotides and Methods of Use

<130> 35718/202438

<150> US 60/166,128

<151> 1999-11-18

<150> US 60/201,837

<151> 2000-05-03

<160> 10

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 824

<212> DNA

<213> Helianthus annuus

<220>

<221> misc\_feature

<222> (0)...(0)

<223> rhoGAP

<221> CDS

<222> (35)...(637)

<400> 1

ttcggcacga gtccaaatcc aatcttcaat cacc atg gct gaa gaa caa ctg ccg  
 55

Met Ala Glu Glu Gln Leu Pro  
 1 5

cct gat caa att aaa ctc att cac aag ctt aat ttg ttc aaa atc aaa  
 103

Pro Asp Gln Ile Lys Leu Ile His Lys Leu Asn Leu Phe Lys Ile Lys  
 10 15 20

ggc aga gat aaa cac aat cgc aaa atc tta cga att gtc gga aaa aac  
 151

Gly Arg Asp Lys His Asn Arg Lys Ile Leu Arg Ile Val Gly Lys Asn  
 25 30 35

ttt cca gct aag agt ttg acc gtt gac ctg ttg aaa aaa tat cta gaa  
 199

Phe Pro Ala Lys Ser Leu Thr Val Asp Leu Leu Lys Lys Tyr Leu Glu  
 40 45 50 55

gtg aaa att ttc ccc aaa ctt gaa cga ccg ttt gtg gtg gtt tac gtt  
 247

Val Lys Ile Phe Pro Lys Leu Glu Arg Pro Phe Val Val Val Tyr Val  
 60 65 70

cac act gat gtt cag aag agc gag aat ttc cct gga ata tcc gtt ctc  
295

His Thr Asp Val Gln Lys Ser Glu Asn Phe Pro Gly Ile Ser Val Leu  
75 80 85

cgg tca gtt tac gac gcg att ccg atg acc gtg aaa caa tat ctt gag  
343

Arg Ser Val Tyr Asp Ala Ile Pro Met Thr Val Lys Gln Tyr Leu Glu  
90 95 100

gcg gtt tac ttt gtt cat ccg gat ctg cag tcc aga att ttt ctg gct  
391

Ala Val Tyr Phe Val His Pro Asp Leu Gln Ser Arg Ile Phe Leu Ala  
105 110 115

aca ttt ggc cgg ctt atc ttc acc gga ggg tta tat gca aag ctg aga  
439

Thr Phe Gly Arg Leu Ile Phe Thr Gly Gly Leu Tyr Ala Lys Leu Arg  
120 125 130 135

ttt gtg agt cga ttg gcg tat ctg tgg gaa cat gtg aaa agg aac gag  
487

Phe Val Ser Arg Leu Ala Tyr Leu Trp Glu His Val Lys Arg Asn Glu  
140 145 150

atc gag atc cca gag ttt gtc tac gat cat gat gag gat ctg gag tac  
535

Ile Glu Ile Pro Glu Phe Val Tyr Asp His Asp Glu Asp Leu Glu Tyr  
155 160 165

cgt ccg atg atg gat tac ggg ata gag agt gac cac gct aga gtt tat  
583

Arg Pro Met Met Asp Tyr Gly Ile Glu Ser Asp His Ala Arg Val Tyr  
170 175 180

gga gcg ccc gcg gtt gat tcc tct gtg gcg gct tat tcc atg agg tgt  
631

Gly Ala Pro Ala Val Asp Ser Ser Val Ala Ala Tyr Ser Met Arg Cys  
185 190 195

atc tca taggggaaat agttgttttt tcttttgttt ttgaaaatag gtgctaaaaag  
687

Ile Ser  
200

aagtgaata tatagtattt agcaatattt cgggtgttgt agtatgttga taacgggctt  
747

ttcttataac attcattgtt ctagttttct ttgttaaaaa ttatttgata aattctttgt  
807

aaaaaaaaaaaa  
824

<210> 2

<211> 201

<212> PRT

<213> Helianthus annuus

<400> 2

Met Ala Glu Glu Gln Leu Pro Pro Asp Gln Ile Lys Leu Ile His Lys  
1 5 10 15

Leu Asn Leu Phe Lys Ile Lys Gly Arg Asp Lys His Asn Arg Lys Ile  
20 25 30

Leu Arg Ile Val Gly Lys Asn Phe Pro Ala Lys Ser Leu Thr Val Asp  
35 40 45

Leu Leu Lys Lys Tyr Leu Glu Val Lys Ile Phe Pro Lys Leu Glu Arg  
50 55 60

Pro Phe Val Val Val Tyr Val His Thr Asp Val Gln Lys Ser Glu Asn

Phe Gly Ala Lys Thr Leu Tyr Ser Val Leu His Gly Val Val Pro Met  
 320 325 330  
 cta gac tca att gta aca gac aaa gac aag ggg ttt cca tta ttc aca  
 1058  
 Leu Asp Ser Ile Val Thr Asp Lys Asp Lys Gly Phe Pro Leu Phe Thr  
 335 340 345  
 tcc ata gat ttg ctt tat aat gaa ggt gtt aat gtt cct tct cct gac  
 1106  
 Ser Ile Asp Leu Leu Tyr Asn Glu Gly Val Asn Val Pro Ser Pro Asp  
 350 355 360  
 aat gga att cta agt gct tta cct aga ctt gtc aaa ggg gct act gat  
 1154  
 Asn Gly Ile Leu Ser Ala Leu Pro Arg Leu Val Lys Gly Ala Thr Asp  
 365 370 375  
 gcc gca aat acc gtt atc aag ttc gag acc ccc gaa acc att gat aga  
 1202  
 Ala Ala Asn Thr Val Ile Lys Phe Glu Thr Pro Glu Thr Ile Asp Arg  
 380 385 390 395  
 gac gca ttc tca tgg ttc cgt gat gaa gag ttc tgc cgg caa atg ctt  
 1250  
 Asp Ala Phe Ser Trp Phe Arg Asp Glu Glu Phe Cys Arg Gln Met Leu  
 400 405 410  
 gcc ggt att aat cct tgt cgc ata caa ttg gtt acg gaa tgg cca ttg  
 1298  
 Ala Gly Ile Asn Pro Cys Arg Ile Gln Leu Val Thr Glu Trp Pro Leu  
 415 420 425  
 atg agt aaa ctg gac cct gaa atc tat gga cca gct gag tca gca att  
 1346  
 Met Ser Lys Leu Asp Pro Glu Ile Tyr Gly Pro Ala Glu Ser Ala Ile  
 430 435 440  
 aca aag gag att gta gag gaa gag att aaa ggt ttc atg act ctt gag  
 1394  
 Thr Lys Glu Ile Val Glu Glu Glu Ile Lys Gly Phe Met Thr Leu Glu  
 445 450 455  
 gag gct tta gca caa aag aag ctg ttt atg ctg gat tat cat gat ctg  
 1442  
 Glu Ala Leu Ala Gln Lys Lys Leu Phe Met Leu Asp Tyr His Asp Leu  
 460 465 470 475  
 ctc ttg cct tat gtt aac aaa acg gag gct gaa ggg aga act ttg tat  
 1490  
 Leu Leu Pro Tyr Val Asn Lys Thr Glu Ala Glu Gly Arg Thr Leu Tyr  
 480 485 490  
 ggt tca aga act tta atg ttc ctt act cct gct gga aca tta agg cca  
 1538  
 Gly Ser Arg Thr Leu Met Phe Leu Thr Pro Ala Gly Thr Leu Arg Pro  
 495 500 505  
 cta gcc att gag ctg act cgc cca cca att gat ggg aaa cca cag tgg  
 1586  
 Leu Ala Ile Glu Leu Thr Arg Pro Pro Ile Asp Gly Lys Pro Gln Trp  
 510 515 520  
 aaa cat gtt tac aca ccc gct tgg gat gct aca ggt gca tgg ctt tgg  
 1634  
 Lys His Val Tyr Thr Pro Ala Trp Asp Ala Thr Gly Ala Trp Leu Trp  
 525 530 535

aag cta gcc aag gct cat gtc ctt gcc cat gat tct agc tat cac caa  
1682  
Lys Leu Ala Lys Ala His Val Leu Ala His Asp Ser Ser Tyr His Gln  
540 545 550 555

ctt gtt agc cat tgg cta aga aca cat tgt gct acc gaa cct tac att  
1730  
Leu Val Ser His Trp Leu Arg Thr His Cys Ala Thr Glu Pro Tyr Ile  
560 565 570

att gct acc aat cgc caa ctc agt caa atg cat cca att cga cga ttt  
1778  
Ile Ala Thr Asn Arg Gln Leu Ser Gln Met His Pro Ile Arg Arg Phe  
575 580 585

cta ctc cct cac ttt cgt tac act atg caa att aat tct cta gct aga  
1826  
Leu Leu Pro His Phe Arg Tyr Thr Met Gln Ile Asn Ser Leu Ala Arg  
590 595 600

ctt tta ctc gtc aat gcc atg ggt atc ata gag tca aca ttt tct cct  
1874  
Leu Leu Leu Val Asn Ala Met Gly Ile Ile Glu Ser Thr Phe Ser Pro  
605 610 615

gga aga tat tgt atg caa att tcc tct gat gca tat gat cag caa tgg  
1922  
Gly Arg Tyr Cys Met Gln Ile Ser Ser Asp Ala Tyr Asp Gln Gln Trp  
620 625 630 635

cgt ttt gat cat gaa gcg ctt ccg gcc gac cta att agc agg ggt atg  
1970  
Arg Phe Asp His Glu Ala Leu Pro Ala Asp Leu Ile Ser Arg Gly Met  
640 645 650

gcg gtt gaa gat cca acc gca cca tat ggt gta aaa cta aca atc gag  
2018  
Ala Val Glu Asp Pro Thr Ala Pro Tyr Gly Val Lys Leu Thr Ile Glu  
655 660 665

gat tac cca tat gca aat gat ggt tta ctc att tat gat acc att aaa  
2066  
Asp Tyr Pro Tyr Ala Asn Asp Gly Leu Leu Ile Tyr Asp Thr Ile Lys  
670 675 680

caa tgg gca act tct tat gtc aac cac tat tac cca cca gcg aat cta  
2114  
Gln Trp Ala Thr Ser Tyr Val Asn His Tyr Tyr Pro Pro Ala Asn Leu  
685 690 695

gtg gaa tct gat gaa gag ctt caa gca tgg tgg aat gaa atc cgt aca  
2162  
Val Glu Ser Asp Glu Glu Leu Gln Ala Trp Trp Asn Glu Ile Arg Thr  
700 705 710 715

gtt ggt cat gga gat aag aaa gat gaa cca tgg tgg cca caa ctc aaa  
2210  
Val Gly His Gly Asp Lys Lys Asp Glu Pro Trp Trp Pro Gln Leu Lys  
720 725 730

acc caa gat gat ttg att gga att gtt tca acc atc ttg tgg gtg acc  
2258  
Thr Gln Asp Asp Leu Ile Gly Ile Val Ser Thr Ile Leu Trp Val Thr  
735 740 745

545                      550                      555                      560  
 Leu Arg Thr His Cys Ala Thr Glu Pro Tyr Ile Ile Ala Thr Asn Arg  
                                  565                      570                      575  
 Gln Leu Ser Gln Met His Pro Ile Arg Arg Phe Leu Leu Pro His Phe  
                                  580                      585                      590  
 Arg Tyr Thr Met Gln Ile Asn Ser Leu Ala Arg Leu Leu Val Asn  
                                  595                      600                      605  
 Ala Met Gly Ile Ile Glu Ser Thr Phe Ser Pro Gly Arg Tyr Cys Met  
                                  610                      615                      620  
 Gln Ile Ser Ser Asp Ala Tyr Asp Gln Gln Trp Arg Phe Asp His Glu  
                                  625                      630                      635  
 Ala Leu Pro Ala Asp Leu Ile Ser Arg Gly Met Ala Val Glu Asp Pro  
                                  645                      650                      655  
 Thr Ala Pro Tyr Gly Val Lys Leu Thr Ile Glu Asp Tyr Pro Tyr Ala  
                                  660                      665                      670  
 Asn Asp Gly Leu Leu Ile Tyr Asp Thr Ile Lys Gln Trp Ala Thr Ser  
                                  675                      680                      685  
 Tyr Val Asn His Tyr Tyr Pro Pro Ala Asn Leu Val Glu Ser Asp Glu  
                                  690                      695                      700  
 Glu Leu Gln Ala Trp Trp Asn Glu Ile Arg Thr Val Gly His Gly Asp  
                                  705                      710                      715  
 Lys Lys Asp Glu Pro Trp Trp Pro Gln Leu Lys Thr Gln Asp Asp Leu  
                                  725                      730                      735  
 Ile Gly Ile Val Ser Thr Ile Leu Trp Val Thr Ser Gly Gln His Ser  
                                  740                      745                      750  
 Ala Val Asn Phe Gly Gln Tyr Asp Phe Ala Gly Tyr Phe Pro Asn Arg  
                                  755                      760                      765  
 Pro Thr Ile Ser Arg Thr Lys Met Pro Asn Glu Asp Pro Thr Asp Glu  
                                  770                      775                      780  
 Glu Trp Gln Ser Phe Ile Lys Arg Pro Glu Asp Ala Leu Leu Lys Cys  
                                  785                      790                      795  
 Phe Pro Ser Gln Ile Gln Ala Thr Lys Val Met Ala Ile Leu Asp Val  
                                  805                      810                      815  
 Leu Ser Ser His Ser Pro Asp Glu Glu Tyr Ile Gly Gly Asn Ile Glu  
                                  820                      825                      830  
 Ala Ala Trp Glu Ala Glu Pro Ala Ile Lys Ala Ala Phe Glu Glu Phe  
                                  835                      840                      845  
 Arg Gly Arg Leu Asn Glu Leu Glu Ala Ile Ile Asp Ser Arg Asn Thr  
                                  850                      855                      860  
 Asp Pro Asn Leu Lys Asn Arg Ser Gly Ala Gly Leu Val Pro Tyr Gln  
                                  865                      870                      875  
 Leu Leu Lys Pro Tyr Ser Glu Lys Gly Val Thr Gly Arg Gly Val Pro  
                                  885                      890                      895  
 Asn Ser Ile Ser Ile  
                                  900

<210> 5  
 <211> 883  
 <212> DNA  
 <213> Helianthus annuus

<220>  
 <221> promoter  
 <222> (1)...(880)  
 <223> lox promoter

<221> misc\_feature  
 <222> (322)...(327)  
 <223> W-box

<221> misc\_feature  
 <222> (722)...(727)  
 <223> G-box

<221> misc\_feature  
 <222> (808)...(811)  
 <223> TATA box

<400> 5  
 agggcacgcg tggcgcacgg accgggctgg gtatctcatt catcttagct cggttttgga  
 60  
 cgtgggttag ttcgttgcg acctcttcca acatagaaca caaaccaca aataagtaca  
 120  
 taaaacccgc tcattttatt agattttatt ctagtccaaa atacgaaaa atcattgtcc  
 180  
 tatatttggtg aaaaatgcta ttttactta tttttccaac aacacatata cagagagggc  
 240  
 aacaatcggg taacaaactc accagagttg tgaaaattat gaggacttct atctgtcatg  
 300  
 caatttttta tatatttctt tttgaccaa acatgtatac atgactaact aaaaatatag  
 360  
 ttgcgagttg gaaaagggtt atacactata actcatattt acacaatatt gccttgaaca  
 420  
 ttattaacta attacacggg gttgaataat tttgataaaa acttttctat gtgttgaggt  
 480  
 atatctgaac tattaaaata aataccatta atacttcaag attataacat gagaacatta  
 540  
 catatattgt gattttatat tataatttaa taattatttt tttttgaaag gcataattta  
 600  
 ataattataa gcgacaatac ttctacgttt atagtactag gtactttttc caaccacaaa  
 660  
 tcaaatgcat tctagccgta gattgtaaat tattaatgca accctgaaca ataatgcata  
 720  
 acacgtgaaa tcaatgcaga aatgtatcat tcttatccga tgttttccca ttaaataaaa  
 780  
 accttaaaat atagcacatt tcctctctat aaatagagct attttttcaa cttccagatc  
 840  
 acacaaaaca agagtgagag tagagtgact aaagaaaacc atg  
 883

<210> 6  
 <211> 1403  
 <212> DNA  
 <213> Helianthus annuus

<220>  
 <221> misc\_feature  
 <222> (0)...(0)  
 <223> ADH cDNA

<221> CDS  
 <222> (74)...(1216)

<400> 6  
 ttcggcacga gccaaaactc acaatttaat ctcatttcaa gaatattctc tctttcaccg  
 60  
 atcaaacaaa agt atg tcg tcg acc act aca ggc caa gtt att cga tgc  
 109

Met	Ser	Ser	Thr	Thr	Thr	Gly	Gln	Val	Ile	Arg	Cys
1				5					10		

aaa gcc gcg gtg acg tgg gaa gcc gga aaa ccg ctg gtg atc gaa gaa  
 157

Lys	Ala	Ala	Val	Thr	Trp	Glu	Ala	Gly	Lys	Pro	Leu	Val	Ile	Glu	Glu
	15						20					25			

gtg gag gtg gcg cca ccg cag aaa atg gaa gtc cgg att aag atc ctc  
 205

Val	Glu	Val	Ala	Pro	Pro	Gln	Lys	Met	Glu	Val	Arg	Ile	Lys	Ile	Leu
	30					35					40				

ttc act tcc ctc tgc cac act gat gtt tac ttc tgg gaa gcc aaa gga  
 253

Phe Thr Ser Leu Cys His Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly

45                      50                      55                      60  
 caa aat cct gta ttc cca aga att tta gga cat gaa gct gga ggg gtt  
 301  
 Gln Asn Pro Val Phe Pro Arg Ile Leu Gly His Glu Ala Gly Gly Val  
                     65                      70                      75  
 gtg gag agt gtt ggg gaa gga gtg act gat ctt cag cca ggg gat cat  
 349  
 Val Glu Ser Val Gly Glu Gly Val Thr Asp Leu Gln Pro Gly Asp His  
                     80                      85                      90  
 gtt ctt ccc gtt ttc acc gga gaa tgc aaa gag tgt gct cac tgt aag  
 397  
 Val Leu Pro Val Phe Thr Gly Glu Cys Lys Glu Cys Ala His Cys Lys  
                     95                      100                      105  
 tcc gaa gag agc aac atg tgt gac ctt ctc agg atc aac acc gac agg  
 445  
 Ser Glu Glu Ser Asn Met Cys Asp Leu Leu Arg Ile Asn Thr Asp Arg  
                     110                      115                      120  
 gga gtc atg ctt cac gat cag aaa tct cga ttc tcg atc aac ggc aaa  
 493  
 Gly Val Met Leu His Asp Gln Lys Ser Arg Phe Ser Ile Asn Gly Lys  
                     125                      130                      135                      140  
 ccc atc ttc cat ttt gtg ggg act tct act ttc agc gag tac acg gtt  
 541  
 Pro Ile Phe His Phe Val Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val  
                     145                      150                      155  
 gtt cat gtt gga tgt ctt gca aag atc aac cct ctt gcc cct ctt gat  
 589  
 Val His Val Gly Cys Leu Ala Lys Ile Asn Pro Leu Ala Pro Leu Asp  
                     160                      165                      170  
 aaa gtt tgt gtt ctc agc tgt ggg atc tcc aca ggg ctg ggt gct act  
 637  
 Lys Val Cys Val Leu Ser Cys Gly Ile Ser Thr Gly Leu Gly Ala Thr  
                     175                      180                      185  
 ttg aat gtt gca aaa ccg aaa aaa ggc tct tcg gtg gcg gtt ttc ggt  
 685  
 Leu Asn Val Ala Lys Pro Lys Lys Gly Ser Ser Val Ala Val Phe Gly  
                     190                      195                      200  
 ctg ggg gca gtg gga ctt gct gct gct gaa ggt gca aga att tct ggg  
 733  
 Leu Gly Ala Val Gly Leu Ala Ala Ala Glu Gly Ala Arg Ile Ser Gly  
                     205                      210                      215                      220  
 gct tca aga atc att ggt gtt gat ctc aat gcc aat aga ttc gag ctt  
 781  
 Ala Ser Arg Ile Ile Gly Val Asp Leu Asn Ala Asn Arg Phe Glu Leu  
                     225                      230                      235  
 gca aag aaa ttt ggg gtt aca gag ttt gtg aac cca aaa gat tat aag  
 829  
 Ala Lys Lys Phe Gly Val Thr Glu Phe Val Asn Pro Lys Asp Tyr Lys  
                     240                      245                      250  
 aag ccg gtg caa gaa gtg att gca gag atg aca aat gga gga gtt gac  
 877  
 Lys Pro Val Gln Glu Val Ile Ala Glu Met Thr Asn Gly Gly Val Asp  
                     255                      260                      265



agg agt gtt gaa tgc act ggt cat att gat gct atg atc tct gct ttt  
 925  
 Arg Ser Val Glu Cys Thr Gly His Ile Asp Ala Met Ile Ser Ala Phe  
 270 275 280

gaa tgt gtt cat gat ggg tgg ggt gtt gct gtt cta gta ggt gtt ccg  
 973  
 Glu Cys Val His Asp Gly Trp Gly Val Ala Val Leu Val Gly Val Pro  
 285 290 295 300

cat aaa gat gcc gtg ttc aag acc agt ccc atg aat ctg ttg aac gaa  
 1021  
 His Lys Asp Ala Val Phe Lys Thr Ser Pro Met Asn Leu Leu Asn Glu  
 305 310 315

agg act ctg aag ggt acc ttc ttt gga aac tat aaa ccg cga tct gat  
 1069  
 Arg Thr Leu Lys Gly Thr Phe Phe Gly Asn Tyr Lys Pro Arg Ser Asp  
 320 325 330

att cct tcg gtt gtc gaa aag tat atg aac aag gaa ctt gag gtg gag  
 1117  
 Ile Pro Ser Val Val Glu Lys Tyr Met Asn Lys Glu Leu Glu Val Glu  
 335 340 345

aag ttc ata aca cat gaa gtg cca ttt tca gag atc aat aag ccc ttt  
 1165  
 Lys Phe Ile Thr His Glu Val Pro Phe Ser Glu Ile Asn Lys Pro Phe  
 350 355 360

gac ttg atg ctt aaa ggt gaa ggt ctt cgt tgc att att cga atg gat  
 1213  
 Asp Leu Met Leu Lys Gly Glu Gly Leu Arg Cys Ile Ile Arg Met Asp  
 365 370 375 380

gcc taaataattt caaactgtgc aagagagagc agtaggagtc gtctattcgt  
 1266  
 Ala

aaagatatat gtgtgtgttc tcgtctctca tcgtcgtaaa tgtgtcctta agatcttggg  
 1326  
 ttgttaattg ttaccataa aagattttga atttgaataa caatagaaat tgatgtctaa  
 1386  
 aaaaaaaaaa aaaaaaa  
 1403

<210> 7  
 <211> 381  
 <212> PRT  
 <213> Helianthus annuus ADH

<400> 7  
 Met Ser Ser Thr Thr Thr Gly Gln Val Ile Arg Cys Lys Ala Ala Val  
 1 5 10 15  
 Thr Trp Glu Ala Gly Lys Pro Leu Val Ile Glu Glu Val Glu Val Ala  
 20 25 30  
 Pro Pro Gln Lys Met Glu Val Arg Ile Lys Ile Leu Phe Thr Ser Leu  
 35 40 45  
 Cys His Thr Asp Val Tyr Phe Trp Glu Ala Lys Gly Gln Asn Pro Val  
 50 55 60  
 Phe Pro Arg Ile Leu Gly His Glu Ala Gly Gly Val Val Glu Ser Val  
 65 70 75 80  
 Gly Glu Gly Val Thr Asp Leu Gln Pro Gly Asp His Val Leu Pro Val  
 85 90 95  
 Phe Thr Gly Glu Cys Lys Glu Cys Ala His Cys Lys Ser Glu Glu Ser  
 100 105 110

Asn Met Cys Asp Leu Leu Arg Ile Asn Thr Asp Arg Gly Val Met Leu  
 115 120 125  
 His Asp Gln Lys Ser Arg Phe Ser Ile Asn Gly Lys Pro Ile Phe His  
 130 135 140  
 Phe Val Gly Thr Ser Thr Phe Ser Glu Tyr Thr Val Val His Val Gly  
 145 150 155 160  
 Cys Leu Ala Lys Ile Asn Pro Leu Ala Pro Leu Asp Lys Val Cys Val  
 165 170 175  
 Leu Ser Cys Gly Ile Ser Thr Gly Leu Gly Ala Thr Leu Asn Val Ala  
 180 185 190  
 Lys Pro Lys Lys Gly Ser Ser Val Ala Val Phe Gly Leu Gly Ala Val  
 195 200 205  
 Gly Leu Ala Ala Ala Glu Gly Ala Arg Ile Ser Gly Ala Ser Arg Ile  
 210 215 220  
 Ile Gly Val Asp Leu Asn Ala Asn Arg Phe Glu Leu Ala Lys Lys Phe  
 225 230 235 240  
 Gly Val Thr Glu Phe Val Asn Pro Lys Asp Tyr Lys Lys Pro Val Gln  
 245 250 255  
 Glu Val Ile Ala Glu Met Thr Asn Gly Gly Val Asp Arg Ser Val Glu  
 260 265 270  
 Cys Thr Gly His Ile Asp Ala Met Ile Ser Ala Phe Glu Cys Val His  
 275 280 285  
 Asp Gly Trp Gly Val Ala Val Leu Val Gly Val Pro His Lys Asp Ala  
 290 295 300  
 Val Phe Lys Thr Ser Pro Met Asn Leu Leu Asn Glu Arg Thr Leu Lys  
 305 310 315 320  
 Gly Thr Phe Phe Gly Asn Tyr Lys Pro Arg Ser Asp Ile Pro Ser Val  
 325 330 335  
 Val Glu Lys Tyr Met Asn Lys Glu Leu Glu Val Glu Lys Phe Ile Thr  
 340 345 350  
 His Glu Val Pro Phe Ser Glu Ile Asn Lys Pro Phe Asp Leu Met Leu  
 355 360 365  
 Lys Gly Glu Gly Leu Arg Cys Ile Ile Arg Met Asp Ala  
 370 375 380

<210> 8  
 <211> 747  
 <212> DNA  
 <213> Helianthus annuus

<220>  
 <221> misc\_feature  
 <222> (0)...(0)  
 <223> SCIP-1 cDNA

<221> CDS  
 <222> (15)...(518)

<400> 8  
 ttccggcacga gcaa atg gcg aac gca agc gat gag ttc aga cta gcg tct  
 50

Met Ala Asn Ala Ser Asp Glu Phe Arg Leu Ala Ser  
 1 5 10

tcc ggc atc gat cat gaa ggc cga cta cca cga aaa tac acc ggt gac  
 98

Ser Gly Ile Asp His Glu Gly Arg Leu Pro Arg Lys Tyr Thr Gly Asp  
 15 20 25

ggt caa ggt aca aaa aaa gac ata tca cca ccg tta gaa tgg tac aac  
 146

Gly Gln Gly Thr Lys Lys Asp Ile Ser Pro Pro Leu Glu Trp Tyr Asn  
 30 35 40

gtt ccg gag ggg aca aaa aca cta gca cta gtg gtg gag gac atc gat  
 194

Val Pro Glu Gly Thr Lys Thr Leu Ala Leu Val Val Glu Asp Ile Asp

```

45              50              55              60
gca ccg gac cca gaa gcg ccg ctg gtt ccg tgg act gtg tgg gtg gtg
242
Ala Pro Asp Pro Glu Ala Pro Leu Val Pro Trp Thr Val Trp Val Val
65              70              75

gtc aat ata cca cct act ttg aag ggg ctc cca gag gga ttt tcc ggg
290
Val Asn Ile Pro Pro Thr Leu Lys Gly Leu Pro Glu Gly Phe Ser Gly
80              85              90

aaa gag ggg gac atg ggt ggc gat tat gct aat gtt aaa gaa gga cat
338
Lys Glu Gly Asp Met Gly Gly Asp Tyr Ala Asn Val Lys Glu Gly His
95              100              105

aat gac ttt aag gtg cct gga tgg cgc gca ccg aag atg ccc tca tcc
386
Asn Asp Phe Lys Val Pro Gly Trp Arg Ala Pro Lys Met Pro Ser Ser
110              115              120

gga cac ccg ttc gag ttt aag ctg tat gcg ttg gat gaa caa gtt gag
434
Gly His Arg Phe Glu Phe Lys Leu Tyr Ala Leu Asp Glu Gln Val Glu
125              130              135              140

ttg ggg aat aag gtg act aag gag aag ttg ctg gag gcg att gat ggc
482
Leu Gly Asn Lys Val Thr Lys Glu Lys Leu Leu Glu Ala Ile Asp Gly
145              150              155

cat gtg gtt ggg gag gct gtt ctg atg gcc gta aat taaattgaga
528
His Val Val Gly Glu Ala Val Leu Met Ala Val Asn
160              165

atggtttata tatatgttag ttgtgtgact tgtgtcatgt gtgatgttct tgttttaacg
588
tattttgaaa cagaagtgac gagagagaga gagtgtttgt tgtgtgtttt tcttgagaga
648
tcgtgaatta attatgctgt ttgcttcaa ggaatcaagc tttataaagt aaaatacaaa
708
tgtaatgctt caaccgagct aaaaaaaaaa aaaaaaaaaa
747

<210> 9
<211> 168
<212> PRT
<213> Helianthus annus SCIP-1

<400> 9
Met Ala Asn Ala Ser Asp Glu Phe Arg Leu Ala Ser Ser Gly Ile Asp
1 5 10 15
His Glu Gly Arg Leu Pro Arg Lys Tyr Thr Gly Asp Gly Gln Gly Thr
20 25 30
Lys Lys Asp Ile Ser Pro Pro Leu Glu Trp Tyr Asn Val Pro Glu Gly
35 40 45
Thr Lys Thr Leu Ala Leu Val Val Glu Asp Ile Asp Ala Pro Asp Pro
50 55 60
Glu Ala Pro Leu Val Pro Trp Thr Val Trp Val Val Val Asn Ile Pro
65 70 75 80
Pro Thr Leu Lys Gly Leu Pro Glu Gly Phe Ser Gly Lys Glu Gly Asp
85 90 95
Met Gly Gly Asp Tyr Ala Asn Val Lys Glu Gly His Asn Asp Phe Lys
100 105 110
Val Pro Gly Trp Arg Ala Pro Lys Met Pro Ser Ser Gly His Arg Phe

```

```

      115      120      125
Glu Phe Lys Leu Tyr Ala Leu Asp Glu Gln Val Glu Leu Gly Asn Lys
      130      135      140
Val Thr Lys Glu Lys Leu Leu Glu Ala Ile Asp Gly His Val Val Gly
      145      150      155      160
Glu Ala Val Leu Met Ala Val Asn
      165

```

```

<210> 10
<211> 513
<212> DNA
<213> Helianthus annuus

```

```

<220>
<221> promoter
<222> (1)...(510)
<223> SCIP promoter

```

```

<221> misc_feature
<222> (364)...(368)
<223> W-box

```

```

<221> misc_feature
<222> (371)...(375)
<223> W-box

```

```

<221> misc_feature
<222> (415)...(420)
<223> G-box

```

```

<400> 10
cttccctatt ttccgtaaca cttgtgcggc aaaggggttg gcagtggta ccgctcggtg
60
ccgaaccact ttccgctgc cactccgggc agcctaaata atgctatata tgtgacattt
120
ttgcactgaa ttctactttt tatttaccat acgcgatgaa aaggcattgg tttttatta
180
tattatattt cagtttctat ttttggacgg caaaaatgaa ttttattaaa agtaaacgaa
240
tttaaaaata ttccgataat tactttttct tttgaatctt gattcggata agttgttacg
300
aattttaaaa cgacaattga ttgaaaatga gtgatgtagc tctttctagc gtaccacgta
360
tctgtcaagt gtcaacatgc tacagcttct caaaactgct agaactctta actacacgtg
420
tccacaaacc cacaaaatcc taaccatcca taacactata agaacttgat caacagatct
480
gtttagtaac aagttattga aggtacaaca atg
513

```

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number  
**WO 01/36464 A3**

(51) International Patent Classification<sup>7</sup>: **C12N 15/29**,  
15/82, 15/11, 9/02, C07K 14/415, A01H 5/00

(21) International Application Number: **PCT/US00/31187**

(22) International Filing Date:  
13 November 2000 (13.11.2000)

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
60/166,128 18 November 1999 (18.11.1999) **US**  
60/201,837 3 May 2000 (03.05.2000) **US**

(71) Applicants (for all designated States except US): **PIONEER HI-BRED INTERNATIONAL, INC.** [US/US];  
800 Capital Square, 400 Locust Street, Des Moines, IA  
50309 (US). **CURAGEN CORPORATION** [US/US];  
555 Long Wharf Drive, 13th Floor, New Haven, CT 06511  
(US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BIDNEY, Dennis**,

**L.** [US/US]; 8385 Plum Drive, Urbandale, IA 50322 (US).  
**DUVICK, Jonathan, P.** [US/US]; 1707 - 38th Street, Des  
Moines, IA 50310 (US). **HENDRICK, Carol** [US/US];  
1707 - 38th Street, Des Moines, IA 50310 (US). **HU, Xu**  
[CA/US]; 4700 - 103rd Street, Urbandale, IA 50322 (US).  
**LU, Guihua** [CN/US]; 4600 - 62nd Street, Urbandale, IA  
50322 (US). **CRASTA, Oswald, R.** [IN/US]; 95-2C Flo-  
rence Road, Branford, CT 06405 (US).


(74) Agents: **WILLIAMSON, Kelly, J. et al.**; Alston & Bird  
LLP, Bank of America Plaza, 101 South Tryon Street, Suite  
4000, Charlotte, NC 28280-4000 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT  
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility  
model), DK, DK (utility model), DM, DZ, EE, EE (utility  
model), ES, FI, FI (utility model), GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK  
(utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian

[Continued on next page]

(54) Title: **SUNFLOWER RhoGAP, LOX, ADH, AND SCIP-1 POLYNUCLEOTIDES AND METHODS OF USE**



(57) Abstract: Methods and compositions for modulating development and defense response are provided. Nucleotide sequences encoding a sunflower rhoGAP, LOX, ADH, and SCIP-1 protein are provided. Nucleotide sequences comprising the LOX and SCIP-1 promoters are also provided. The sequences can be used in expression cassettes for modulating development, developmental pathways, and the plant defense response. Transformed plants, plant cells, tissues, and seed are also provided.

WO 01/36464 A3



patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:  
18 April 2002

**Published:**

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## INTERNATIONAL SEARCH REPORT

In International Application No

PCT/US 00/31187

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/29 C12N15/82 C12N15/11 C12N9/02 C07K14/415  
A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! Entry/Acc.no. 081806, 1 November 1998 (1998-11-01) KOETTER, P. ET AL.: "HYPOTHETICAL 23.6 KDA PROTEIN." XP002174683 the whole document</p>	1-15
X	<p>DATABASE EMBL 'Online! Entry/Acc.no. AW031283, 17 September 1999 (1999-09-17) ALCALA, J. ET AL.: "EST274658 tomato callus, TAMU Lycopersicon esculentum cDNA clone cLEC35A6 similar to A. thaliana putative protein, mRNA sequence." XP002174684 the whole document</p>	1-15

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

3 January 2002

Date of mailing of the international search report

01 02 2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Smalt, R

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 00/31187

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ROYO JOAQUIN ET AL: "Characterization of three potato lipoxygenases with distinct enzymatic activities and different organ-specific and wound-regulated expression patterns." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 35, 1996, pages 21012-21019, XP002186561 ISSN: 0021-9258 the whole document</p>	1-15
X	<p>DATABASE EMBL 'Online! Entry/Acc.no. Q9M042, 1 October 2000 (2000-10-01) BEVAN, M ET AL.: "HYPOTHETICAL 17.8 KDA PROTEIN." XP002186562 the whole document</p>	1-15
X	<p>DATABASE EMBI 'Online! Entry/Acc.no. AI032451, 24 June 1998 (1998-06-24) MICHALOWSKI, C.B. ET AL.: "R6-B5SK Ice plant Lambda Uni-Zap XR expression library, root, 6h NaCl treatment Mesembryanthemum crystallinum cDNA clone R6-B5 5', mRNA sequence." XP002186563 the whole document</p>	2
X	<p>ALMOGUERA D ET AL: "DEVELOPMENT AND ENVIRONMENT CONCURRENT EXPRESSION OF SUNFLOWER DRY-SEED-STORED LOW-MOLECULAR-WEIGHT HEAT-SHOCK PROTEIN AND LEA MRNAS" PLANT MOLECULAR BIOLOGY, NIJHOFF PUBLISHERS, DORDRECHT, NL, vol. 19, no. 5, 1992, pages 781-792, XP000944118 ISSN: 0167-4412 the whole document</p>	17
X	<p>DATABASE EMBL 'Online! Entry/Acc.no. AU034024, 28 October 1998 (1998-10-28) MORIO, T. ET AL.: "Dictyostelium discoideum slug cDNA, clone SLB818." XP002186564 the whole document</p>	17

-/--



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 00/31187

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WINGE ET AL: "Cloning and characterization of rac-like cDNAs from Arabidopsis thaliana" PLANT MOLECULAR BIOLOGY, NIJHOFF PUBLISHERS, DORDRECHT, NL, vol. 35, 1997, pages 483-495, XP002131349 ISSN: 0167-4412 the whole document</p>	
A	<p>TAKAI Y ET AL: "Rho as a regulator of the cytoskeleton" TIBS TRENDS IN BIOCHEMICAL SCIENCES, ELSEVIER PUBLICATION, CAMBRIDGE, EN, vol. 20, no. 6, June 1995 (1995-06), pages 227-231, XP004222365 ISSN: 0968-0004 the whole document</p>	
A	<p>WO 99 04013 A (FALAK IGOR ; NAZARIAN KATHERINE A P (CA); PIONEER HI BRED INT (US);) 28 January 1999 (1999-01-28) the whole document</p>	
A	<p>GENTZBITTEL I ET AL: "CLONING OF MOLECULAR MARKERS FOR DISEASE RESISTANCE IN SUNFLOWER, HELIANTHUS ANNUUS L" THEORETICAL AND APPLIED GENETICS, SPRINGER, BERLIN, DE, vol. 96, no. 3/04, March 1998 (1998-03), pages 519-525, XP001002152 ISSN: 0040-5752 the whole document</p>	
P, X	<p>EP 1 033 405 A (CERES INC) 6 September 2000 (2000-09-06) * see seq.ID's 54103 and 54105 *</p>	<p>1-5, 14, 15</p>

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 00/31187

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:  
1-24
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-6,8-16, all partially

Sunflower rho-GTPase activating protein according to seq.ID.2 and homologues with at least 70% sequence identity thereto, nucleic acids encoding said proteins and homologues thereof having at least 60% sequence identity to seq.ID.1, optionally functionally linked to a promoter, cell or transformed plant or seeds thereof having a genome which has incorporated therein said nucleic acid linked to a promoter, and method for enhancing a plant defence response by incorporating into its genome said nucleic acid sequence linked to a heterologous promoter.

2. Claims: 1-24, all partially

Sunflower lipoxigenase protein according to seq.ID.4 and homologues with at least 60% sequence identity thereto, nucleic acids encoding said proteins and homologues thereof having at least 60% sequence identity to seq.ID.3, optionally functionally linked to a promoter, cell or transformed plant or seeds thereof having a genome which has incorporated therein said nucleic acid linked to a promoter, and method for enhancing a plant defence response by incorporating into its genome said nucleic acid sequence linked to a heterologous promoter.

Also an isolated nucleic acid comprising the promoter sequence of said gene according to seq.ID.5 or homologues thereof which hybridize thereto under stringent conditions, optionally linked to a nucleic acid sequence to be expressed, expression vector comprising said nucleic acid, cell or plant having said nucleic acid linked to said sequence to be expressed stably incorporated into its genome, and method of regulating the expression of a nucleic acid sequence using said promoter.

3. Claims: 1-6,8-16, all partially

Sunflower alcohol dehydrogenase protein according to seq.ID.7 and homologues with at least 90% sequence identity thereto, nucleic acids encoding said proteins and homologues thereof having at least 60% sequence identity to seq.ID.6, optionally functionally linked to a promoter, cell or transformed plant or seeds thereof having a genome which has incorporated therein said nucleic acid linked to a promoter, and method for enhancing a plant defence response by incorporating into its genome said nucleic acid sequence linked to a heterologous promoter.

4. Claims: 1-24, all partially

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Sunflower Sclerotinia-inducible protein-1 protein according to seq.ID.9 and homologues with at least 60% sequence identity thereto, nucleic acids encoding said proteins and homologues thereof having at least 60% sequence identity to seq.ID.8, optionally functionally linked to a promoter, cell or transformed plant or seeds thereof having a genome which has incorporated therein said nucleic acid linked to a promoter, and method for enhancing a plant defence response by incorporating into its genome said nucleic acid sequence linked to a heterologous promoter.

Also an isolated nucleic acid comprising the promoter sequence of said gene according to seq.ID.10 or homologues thereof which hybridize thereto under stringent conditions, optionally linked to a nucleic acid sequence to be expressed, expression vector comprising said nucleic acid, cell or plant having said nucleic acid linked to said sequence to be expressed stably incorporated into its genome, and method of regulating the expression of a nucleic acid sequence using said promoter.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/31187

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9904013	A	28-01-1999	AU	8411198 A	10-02-1999
			WO	9904013 A2	28-01-1999
EP 1033405	A	06-09-2000	EP	1033405 A2	06-09-2000

**THIS PAGE BLANK (USPTO)**